

11. REFERENCIAS

1. Agosti, D. (2000). *Ants: standard methods for measuring and monitoring biodiversity*. Michigan: Smithsonian Institution Press.
2. Andersen, A. (1997). Using Ants as bioindicators: Multiscale Issues in Ant Community Ecology. *Ecology and Society* .
3. Andersen, A. N. (1997). Functional Groups and patterns of organization in North American ant communities: a comparison with Australia. *Journal of Biogeography* , 433-460.
4. Anderssen, A. N. (1995). A classification of Australian ant communities, based on functional groups which parallel plant-life forms in relation to stress and disturbance . *Journal of Biogeography* , 15-29 .
5. Badano, E. I. (2010). RELACIONES ENTRE EL ESTADO DE SALUD DEL ECOSISTEMA DEL PARQUE ESTATAL FLOR DEL BOSQUE Y FACTORES DE CAMBIO GLOBAL . Proyecto Fondo Mixto CONACYT-Gobierno del Estado de Puebla , 1-28.
6. Badano, E. I. (2011). Emulating nurse plants to restore oak forests. *Ecological Engineering* , 1244-1248.
7. Badano, E. I. (2012). Conservation value of a natural protected area in the state of Puebla, Mexico. *Revista Mexicana de Biodiversidad* , 834-846.
8. Burbidge, A. H. (1992). Ants as indicators of disturbance at Yanchep National Park, Western Australia. *Journal of the Royal Society of Western Australia* , 89-95.

9. Callaway, R. (1997). Competition and facilitation: a synthetic approach to interactions in plant communities. *Ecology* , 1958–1965.
10. Cano, J. F. (2012). EFFECTS OF BURIAL DEPTH ON SEED GERMINATION AND SEEDLING EMERGENCE OF MEXICAN OAKS: A GLASSHOUSE EXPERIMENT. *Archives of Biological Science Belgrade* , 1543-1554.
11. Carvalho, K. S. (1999). Forest fragmentation in central Amazonia and its effects on litter-dwelling ants . *Biological Conservation* , 151-157.
12. Casanova, L. R. (2014). Biodiversidad de hormigas en México. *Revista Mexicana de Biodiversidad* , 92-98.
13. Cerdá, X. (1998). The role of competition by dominants and temperature in the foraging of subordinate species in Mediterranean ant communities . *Oecologia* , 404-412.
14. Charnley, S. (2010). Mitigating Climate Change Through Small-Scale Forestry in the USA: Opportunities and Challenges . *Small-scale Forestry* , 445–462 .
15. CONAFOR. (2009). Comisión Nacional Forestal . Recuperado el 5 de Noviembre de 2015, de Restauración de ecosistemas forestales : <http://www.conafor.gob.mx:8080/documentos/docs/7/579Restauración%20de%20ecosistemas%20forestales.pdf>
16. CONAFOR. (24 de Febrero de 2015). Servicios Ambientales. Recuperado el 9 de Noviembre de 2015, de SEMARNAT: <http://www.conafor.gob.mx/web/temas-forestales/servicios-ambientales/>
17. Cuautle, M. (2011). RELACIONES ENTRE EL ESTADO DE SALUD DEL ECOSISTEMA DEL PARQUE ESTATAL FLOR DEL BOSQUE Y

FACTORES DE CAMBIO GLOBAL. Gobierno del Estado de Puebla,
Secretaria de Sustentabilidad Ambiental y Ordenamiento Territorial .

18. Cuautle, M. (2015). Comparison of Ant community Diversity and Functional Group Composition Associated to Land Use Change in a Seasonally Dry Oak Forest in Mexico . *Neotropical Entomology* .
19. Dolek, M. (2008). Ants on oaks: effects of forest structure on species composition . *Journal of Insect Conservation* , 367–375 .
20. Ellison, A. M. (2007). Rapid Inventory of the Ant Assemblage in a Temperate Hardwood Forest: Species Composition and Assessment of Sampling Methods. *Community and Ecosystem Ecology* , 766-775.
21. Fernández, F. (2003). Introducción a las Hormigas de la región Neotropical. Bogotá, Colombia : Instituto de Investigación de Recursos Biológicos Alexander von Humboldt .
22. Fernández, P. R. (2001). LAS HORMIGAS DEL SUELO EN MÉXICO: DIVERSIDAD, DISTRIBUCIÓN E IMPORTANCIA (HYMENOPTERA: FORMICIDAE) . *Acta Zoológica Mexicana* , 189-238.
23. Gomes, E. (Septiembre de 2014). Ant assemblages (Hymenoptera: Formicidae) in three different stages of forest regeneration in a fragment of Atlantic Forest in Sergipe, Brazil. . *Sociobiology* , 250-257.
24. Jackson, R. B. (2008). Protecting climate with forests . *Environmental Research Letters* , 1-5.
25. Jianzhong, Y. (2005). Land cover changes based on plant successions: Deforestation, rehabilitation and degeneration of forest in the upper Dadu River watershed . *Science in China* , 48 (12), 2214—2230 .

26. Liancourt, P. (2005). Stress tolerance and competitive-response ability determine the outcome of biotic interactions. *Ecology* , 1611–1618.
27. López-Barrera, F. (2001). Influence of litter on emergence and early growth of *Quercus rugosa*: a laboratory study. *Boletín de la sociedad Botánica de México* , 59-70.
28. Lortie, C. (2004). Rethinking plant community theory. *Oikos* , 433-438.
29. Luna-José, A. D. (2003). LOS USOS NO LEÑOSOS DE LOS ENCINOS EN MÉXICO. *Boletín de la Sociedad Botánica de México* , 107-117.
30. Majer, J. D. (2007). Ants (Hymenoptera: Formicidae) pass the bioindicator scorecard . *Myrmecological News* , 69-76.
31. Padilla, F. M. (2006). The role of nurse plants in the restoration of degraded environments. *Frontiers in ecology and environment* , 196-202.
32. Pais, M. P. (2010). Arthropod Recolonization in the Restoration of a Semideciduous Forest in Southeastern Brazil. *Neotropical Entomology* , 39 (2), 198-206.
33. Parra, V. (2003). Listado Microfaunístico del Parque Zoológico Africam Safari. *Elementos: ciencia y cultura* , 49-51.
34. Peck, S. L. (1998). Using Ant Species (Hymenoptera: Formicidae) as a Biological Indicator of Agroecosystem Condition . *Community and Ecosystem Ecology* , 1102-1110.
35. R. Ribas, C. (2012). Ants as Indicators in Brazil: A Review with Suggestions to Improve the Use of Ants in Environmental Monitoring Programs. *Hindawi Publishing Corporation* , 2-22.

36. Rico-Gray, V. (1993). Use of Plant-Delivered Food Resources by Ants in the Dry Tropical Lowlands of Coastal Veracruz, Mexico. *Biotropica* , 301-315.
37. Rojas, P. (1996). Formicidae (Himenoptera), Biodiversidad, taxonomía y biogeografía de artrópodos de México: hacia una síntesis de su conocimiento. *Acta Zoologica Mexicana* , 483-500.
38. Santamaría, L. R. (2013). Insectos Fitófagos en las Especies de Encino del Parque Africam Safari en Tecali de Herrera, Puebla. Estado de México : Universidad Autónoma de Chilpancingo.
39. Suguituru, S. S. (2011). Ant community richness and composition across a gradient from Eucalyptus plantations to secondary Atlantic Forest . *Biota Neotropica* , 369-376.
40. Sulea, C. (2013). Assessing the Importance of Biotic Factors in Tree Development. *Bulletin UASMV* , 317-320.
41. Torres, A. B. (2013). Potential for Climate Change Mitigation in Degraded Forests: A Study from La Primavera, Mexico . *Forest* , 1032-1054 .
42. Valencia-A, S. (2004). Diversidad del Género Quercus (Fagaceae) en Mexico. *Boletín de la Sociedad Botánica de México* (75), 33-53.
43. Verzero Villalba, F. (2014). Grupos funcionales dominantes de hormigas (Hymenoptera: Formicidae) en pastizales naturales con y sin pastoreo del noroeste de Buenos Aires, Argentina. *Revista de la Facultad de Agronomía* , 107-113.

