

## Bibliografía

- Aapola U, Shibuya K, Scott HS, Ollila J, Vihtinen M, Heino M, Shintani A, Kawasaki K, Minoshima S, Krohn K, Antonarakis SE, Shimizu N, Kudoh J, Peterson P. Isolation and initial characterization of a novel zinc finger gene, DNMT3L, on 21q22.3, related to the cytosine-5-methyltransferase 3 gene family. *Genomics* 2000;65:293-8.
- Aapola U, Lyle R, Krohn L, Antonarakis SE, Peterson P. Isolation and initial characterization of the mouse Dnmt3l gene. *Cytogenet. Cell Genet.* 2001;92:122-6.
- Aasland R, Gibson T, Stewart AF. The PHD finger: implications for chromatin-mediated transcripcional regulation. *TIBS* 1995;20:56-9.
- ACS Gentuzumab ozogamicin. *American Cancer Society: Cancer Drug Guide*. Disponible en: [http://our.cancer.org/docroot/CDG/content/CDG\\_gemtuzumab\\_ozogamicin.asp](http://our.cancer.org/docroot/CDG/content/CDG_gemtuzumab_ozogamicin.asp). Accesado el 10 junio 2008.
- Abdolmaleky HM, Cheng KH, Faraone SV, Shafa R, Stone W, Glatt SJ, Tsuang MT. Hypomethylation of MB-COMT promoter is a major risk factor for schizophrenia and bipolar disorder. *Hum. Mol. Genet.* 2006;15:3132-45.
- Abumrad NA. The gene-nutrient-gene loop. *Curr. Opin. Clin. Nutr. Metab. Care* 2001;4:207-10.
- Adamietz P, Rudolph A. ADP-ribosylation of nuclear proteins in vivo. Identification of histone H2B as a major acceptor for mono- and poly(ADP-ribose) in dimethyl sulphate-treated hepatoma AH 7974 cells. *J. Biol. Chem.* 1984;259:6841-6.
- Agrelo R, Cheng WH, Setien F, Ropero S, Espada J, Fraga MF, Herranz M, Paz MF, Sanchez-Cespedes M, Artiga MJ, Guerrero D, Castells A, von Kobbe C, Bohr VA, Esteller M. Epigenetic inactivation of the premature aging Werner syndrome gene in human cancer. *Proc. Natl. Acad. Sci. USA* 2006;103:8822-7.
- Ahmad K, Henikoff S. The histone variant H3.3 marks active chromatin by replication independent nucleosome assembly. *Mol. Cell* 2002;9:1191-200.
- Akhtar A, Zink D, Becker PB. Chromodomains are protein-RNA interaction modules. *Nature* 2000;407:405-9.
- Algeciras-Schimmler A, Barnhart B, Peter ME. Apoptosis dependent and independent function caspases. *Eurekah Bioscience Collection 2005*. Disponible en: <http://www.ncbi.nlm.nih.gov/books/bv.fcgi?rid=eurekah>. Accesado el 10 de junio 2008.
- Allard S, Utley RT, Savard J, Clarke A, Grant P, Brandl CJ, Pillus L, Workman JL, Cote J. NuA4, an essential transcription adaptor/histone H4 acetyltransferase complex containing Esa1p and the ATM-related cofactor Tra1p. *EMBO J.* 1999;18:5108-19.
- Allfrey V, Faulkner RM, Mirsky AE. Acetylation and methylation of histones and their possible role in the regulation of RNA synthesis. *Proc. Natl. Acad. Sci.* 1964;51:786-94.
- Allis CD, Bowen JK, Abraham GN, Glover CV, Gorovsky MA. Proteolytic processing of histone H3 in chromatin: a physiologically regulated event in *Tetrahymena micronuclei*. *Cell* 1980;20:55-64.
- Allshire RC, Nimmo ER, Ekwall K, Javerzat JP, Craston G. Mutations derepressing silent centromeric domains in fission yeast disrupt chromosome segregation. *Genes Dev.* 1995;9:218-233.
- ALSPAC Avon Longitudinal Study of Parents and Children. Disponible en: <http://www.alspac.bris.ac.uk/>. Consultado el 10 de junio 2008.
- Alvarez-Gonzalez R, Althaus FR. Poly(ADP-ribose) catabolism in mammalian cells exposed to DNA-damaging agents. *Mutat. Res.* 1989;218:67-74.
- Alvarez-Gonzalez R, Jacobson MK. Characterization of polymers of adenosine diphosphate ribose generated in vitro and in vivo. *Biochemistry* 1987;26:3218-24.
- Ame JC, Jacobson EL, Jacobson MK. Molecular heterogeneity and regulation of poly(ADP-ribose) glycohydrolase. *Mol. Cell Biochem.* 1999a; 193:75-81.

- Ame JC, Rolli V, Schreiber V, Niedergang C, Apiou F, Decker P, Muller S, Höger T, Ménissier-de Murcia J, de Murcia G. PARP-2, a novel mammalian DNA damage-dependent poly(ADP-ribose) polymerase. *J. Biol. Chem.* 1999b; 274:17860-8.
- Ame JC, Spenlehauer C, de Murcia G. The PARP superfamily. *Bioessays* 2004;26:882-93.
- Amir RE, Zoghbi HY. Rett syndrome: methyl-CpG-binding protein 2 mutations and phenotype-genotype correlations. *Am. J. Med. Genet.* 2000;97:147-52.
- An W, Kim J, Roeder RG. Ordered cooperative functions of PRMT1, p300, and CARM1 in transcriptional activation by p53. *Cell* 2004;117:735-48.
- Angrand PO, Apiou F, Stewart AF, Dutrillaux B, Losson R, Chambon P. NSD3, a new SET domain-containing gene, maps to 8p12 and is amplified in human breast cancer. *Genomics* 2001;74:79-88.
- Antequera F, Bird A. Number of CpG islands and genes in human and mouse. *Proc. Natl. Acad. Sci.* 1993;90:11995-9.
- Antun FT, Burnett GB, Cooper AJ, Daly RJ, Smythies JR, Zealley AK. The effects of L-methionine (without MAOI) in schizophrenia. *J. Psychiatr. Res.* 1971;8:63-71.
- Aoki A, Suetake I, Miyagawa J, Fujio T, Chijiwa T, Sasaki H, Tajima S. Enzymatic properties of de novo-type mouse DNA (cytosine-5) methyltransferases. *Nucleic Acids Res.* 2001;29:3506-12.
- Aoki K, Meng G, Suzuki K, Takashi T, Kameoka Y, Nakahara K, Ishida R, Kasai M. RP58 associates with condensed chromatin and mediates a sequence-specific transcriptional repression. *J. Biol. Chem.* 1998;273:26698-704.
- ARGENBIO Consejo Argentino para la Información y el Desarrollo de la Biotecnología. Glosario. Disponible en: <http://www.argenbio.org/h/glosario/>. Accesado el 18 de mayo 2008.
- Artursson P, Palm K, Luthman K. Caco-2 monolayers in experimental and theoretical predictions of drug transport. *Adv. Drug Deliv. Rev.* 2001; 46:27-43.
- Aubin R, Frechette A, de Murcia G, Malouin F, Lord A, Mandel P, Poirier GG. Nucleosomal poly(ADP-ribose) polymerase: properties and relaxation of the chromatin structure. *Princesa Takamatsu Symp.* 1983;13:83-91.
- Augustin A, Spenlehauer C, Dumond, H., Ménissier-de Murcia, J., Piel, M., Schmit, A. C., Apiou, F., Vonesch, J. L., Kock, M., Bornens, M., de Murcia, G. PARP-3 localizes preferentially to the daughter centriole and interferes with the G1/S cell cycle progression. *J. Cell Sci.* 2003;116:1551-62.
- Baarends WM, Hoogerbrugge JW, Roest HP, Ooms M, Vreeburg J, Hoeijmakers JH, Grootegoed JA. Histone ubiquitination and chromatin remodelling in mouse spermatogenesis. *Dev. Biol.* 1999;207:322-33.
- Bach-Knudsen KE, Serena A, Canibe N, Juntunen KS. New insight into butyrate metabolism. *Proc. Nutr. Soc.* 2003;62:81-6.
- Bachman KE, Rountree MR, Baylin SB. Dnmt3a and Dnmt3b are transcriptional repressors that exhibit unique localization properties to heterochromatin. *J. Biol. Chem.* 2001;276:32282-7.
- Bailey D, O'Hare P. Herpes simplex virus 1 ICP0 co-localizes with SUMO-specific protease. *J. Gen. Virol.* 2002;83:2951-64.
- Ballard TD, Wolff J, Griffin JB, Stanley JS, Calcar SV, Zempleni J. Biotinidase catalyzes debiotinylation of histones. *Eur. J. Nutr.* 2002;41:78-84.
- Ballestar E, Wolffe AP. Methyl-CpG-binding proteins. Targeting specific gene expression. *Eur. J. Biochem.* 2001;268:1-6.
- Bannister AJ, Zegerman P, Partridge JF, Miska EA, Thomas JO, Allshire RC, Kouzarides T. Selective recognition of methylated lysine 9 on histone H3 by HP1 chromo domain. *Nature* 2001;410:120-4.
- Bannister AJ, Schneider R, Kouzarides T. Histone methylation: dynamic or static? *Cell* 2002;109:801-6.
- Bantignies F, Goodman RH, Smolik SM. Functional interaction between the coactivator Drosophila CREB-binding protein and ASH1, a member of the trithorax group of chromatin modifier. *Mol. Cell Biol.* 2000;20:9317-30.
- Barker DJ. The fetal and infant origins of adult disease. *BMJ* 1990;301:1111.

- Barker DJ, Gluckman PD, Godfrey KM, Harding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult life. *Lancet* 1993;341:938-41.
- Barlow DP. Gametic imprinting in mammals. *Science* 1995;270:1610-13.
- Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, Wei G, Chepelev I, Zhao K. High-resolution profiling of histone methylations in the human genome. *Cell* 2007;129:823-37.
- Barsoum J, Varshavsky A. Preferential localization of variant nucleosomes near the 5'-end of the mouse dihydrofolate reductase gene. *J. Biol. Chem.* 1985;260:7688-97.
- Bassett SS, Avramopoulos D, FallinD. Evidence for parent of origin effect in late-onset Alzheimer disease. *Am. J. Med. Genet.* 2002;114:679-86.
- Bauer UM, Daujar S, Nielsen SJ, Nightingale K, Kouzarides T. Methylation at arginine 17 of histone H3 is linked to gene activation. *EMBO Rep.* 2001;3:39-44.
- Baumbusch LO, Thorstensen T, Krauss V, Fischer A, Naumann K, Assalkhou R, Schulz1, Reuter IG, and Aalen RB. The Arabidopsis thaliana genome contains at least 29 active genes encoding SET domain proteins that can be assigned to four evolutionary conserved classes. *Nucleic Acids Res.* 2001;29:4319-33.
- Bayer P, Arndt A, Metzger S, Mahajan R, Melchior F, Jaenicke R, Becker J. Structure determination of the small ubiquitin-related modifier SUMO-1. *J. Mol. Biol.* 1998;280:275-86.
- Bayerlein K, Hillemacher T, Reulbach U, Mugele B, Sperling W, Kornhuber J, Bleich S. Alcoholism associated hyperhomocysteinemia and previous withdrawal seizures. *Biol. Psychiatry* 2005;57:1590-3.
- Baylin S, Bestor TH. Altered methylation patterns in cancer cell genomes: cause or consequence? *Cancer Cell* 2002;1:299-305.
- Bedford MT, Richard S. Arginine methylation an emerging regulator of protein function. *Mol. Cell* 2005;18:263-72.
- Beiby J, Ingram D, Hahnel R, Rossi E. Reduced breast cancer risk with increasing serum folate in a case-control study of the C677T genotype of the methylenetetrahydrofolate reductase gene. *Eur. J. Cancer* 2004;40:1250-4.
- Beisel C, Imhof A, Greene J, Kremmer E, Sauer F. Histone methylation of the Drosophila epigenetic transcriptional regulator Ash1. *Nature* 2002;419:857-62.
- Bernier-Villamor V, Sampson DA, Matunis MJ, Lima CD. Structural basis for E2-mediated SUMO conjugation revealed by a complex between ubiquitin-conjugating enzyme Ubc9 and RanGAP1. *Cell* 2002;108:345-56.
- Bernstein BE, Kamal M, Lindblad-Toh K, Bekiranov S, Bailey DK, Huebert DJ, McMahon S, Karlsson EK, Kulbokas EJ 3rd, Gingeras TR, Schreiber SL, Lander ES. Genome maps and comparative analysis of histone modifications in human and mice. *Cell* 2005;120:169-81.
- Bertos NR, Wang AH, Yang XJ. Class II histone deacetylases: structure function and regulation. *Biochem. Cell Biol.* 2001;79:243-52.
- Bestor TH. The DNA methyltransferases of mammals. *Hum. Mol. Genet.* 2000;9:2395-402.
- Bhave MR, Wilson MJ, Poirier LA. c-H-ras and c-K-ras gene hypomethylation in the livers and hepatomas of rats fed methyl-deficient, aminoacid-definer diets. *Carcinogenesis* 1988;9:343-8.
- Biade S, Stobbe CC, Boyd JT, Chapman JD. Chemical agents that promote chromatin compaction radiosensitize tumour cells. *Int. J. Radiat. Biol.* 2001;77:1033-42.
- Bielawski DM, Zaher FM, Svinarich DM, Abel EL. Paternal alcohol exposure affects sperm cytosine methyltransferase messenger RNA levels. *Alcohol Clin. Exp. Res.* 2002;26:347-51.
- Binisziewicz D, Gribnau J, Ramsahoye B, Gaudet F, Eggan K, Humpherys D, Mastrangelo MA, Jun Z, Walter J, Jaenisch R. DNMT1 overexpression causes genomic hypermethylation, loss of imprinting, and embryonic lethality. *Mol. Cell. Biol.* 2002;22:2124-35.
- Bird A. Molecular biology: methylation talk between histones and DNA. *Science* 2001; 294:2113-5.
- Blander G, Guarente L. The Sir2 family of protein deacetylases. *Annu. Rev. Biochem.* 2004;73:417-435.
- Bleich S, Degner D, Wiltfang J, Maler JM, Niedmann P, Cohrs S, Mangholz A, Porzig J, Sprung R, Rüther E, Kornhuber J. Elevated homocysteine levels in alcohol withdrawal. *Alcohol Alcoholism* 2000a;35:351-4.

- Bleich S, Degner D, Bandelow B, von Ahsen N, Rüther E, Kornhuber J. Homocysteine is a predictor of alcohol withdrawal seizures. *Neuroreport* 2000b;11:2749-52.
- Bleich S, Bandelow B, Javaheripour K, Müller A, Degner D, Wilhelm J, Havemann-Reinecke U, Sperling W, Rüther E, Kornhuber J. Hyperhomocysteinemia as a new risk factor for brain shrinkage in patients with alcoholism. *Neurosci. Lett.* 2003;335:179-82.
- Bleich S, Bayerlein K, Reulbach U, Hillemacher T, Bonsch D, Mugele B, Kornhuber J, Sperling W. Homocysteine levels in patients classified according to Lesch's typology. *Alcohol Alcoholism* 2004;39:493-8.
- Bleich S, Carl M, Bayerlein K, Reulbach U, Biermann T, Hillemacher T, Bonsch D, Kornhuber J. Evidence of elevated homocysteine levels in alcoholism. The Franconian Alcoholism Research Studies (FARS). *Alcohol Clin. Exp. Res.* 2005;29:334-6.
- Bleich S, Lenz B, Ziegenbein M, Beutler S, Frieling H, Kornhuber J, Bonsch D. Epigenetic DNA hypermethylation of the HERP gene promoter induces down-regulation of its mRNA expression in patients with alcohol dependence. *Alcohol Clin. Exp. Res.* 2006;30:587-91.
- Blondeau B, Avril I, Duchene B, Breant B. Endocrine pancreas development is altered in foetuses from rats previously showing intra-uterine growth retardation in response to malnutrition. *Diabetologia* 2002;45:394-401.
- Bohm L, Schneeweiss FA, Sharan RN, Feinendegen LE. Influence of histone acetylation on the modification of cytoplasmic and nuclear proteins by ADP-ribosylation in response to free radicals. *Biochim. Biophys. Acta* 1997;1334:149-54.
- Bolden JE, Peart MJ, Johnstone RW. Anticancer activities of histone deacetylase inhibitors. *Nat. Rev. Drug Discov.* 2006;5:769-84.
- Boloker J, Gertz SJ, Simmons RA. Gestational diabetes leads to the development of diabetes in adulthood in the rat. *Diabetes* 2002;51:1499-506.
- Boltze C, Schneider-Stock R, Quednow C, Hinze R, Mawrin C, Hribaschek A, Roessner A, Hoang-Vu C. Silencing of the maspin gene by promoter hypermethylation in thyroid cancer. *Int. J. Mol. Med.* 2003;12:479-84.
- Bonfils C, Beaulieu N, Chan E, Cotton-Montpetit J, MacLeod AR. Characterization of the human DNA methyltransferase splice variant Dnmt1b. *J. Biol. Chem.* 2000;275:10754-60.
- Bonicalzi ME, Haince JF, Droit A, Poirier GG. Regulation of poly(ADP-ribose) metabolism by poly(ADP-ribose) glycohydrolase: where and when? *Cell Mol. Life Sci.* 2005;62:739-50.
- Bonitaiti-Pellie C. Methodological aspects of investigating gene-nutrient interactions. *Eur. J. Cancer Prev.* 2002;11:69-74.
- Borrow J, Stanton VPJ, Andresen JM, Becher R, Behm FG, Chaganti RS, Civin CI, Disteche C, Dubé I, Frischauf AM, Horsman D, Mitelman F, Volinia S, Watmore AE, Housman DE. The translocation t(8;16)(p11;p13) of acute myeloid leukaemia fuses a putative acetyltransferase to the CREB-binding protein. *Nature Genet.* 1996;14:33-41.
- Bottomley MJ. Structures of protein domains that create or recognize histone modifications. *EMBO Rep.* 2004;5:464-9.
- Bouillet P, Strasser A. BH3-only proteins- evolutionarily conserved proapoptotic Bcl-2 family members essential for initiating programmed cell death. *J. Cell. Sci.* 2002;115:1567-74.
- Boulanger MC, Liang C, Russell RS, Lin R, Bedford MT, Wainberg MA, Richard S. Methylation of Tat by PRMT6 regulates human immunodeficiency virus type 1 gene expression. *J. Virol.* 2005;79:124-31.
- Boulikas T. Relation between carcinogenesis, chromatin structure and poly(ADP-ribosylation). *Anticancer Res.* 1991;11:489-527.
- Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH. Dnmt3L and the establishment of maternal genomic imprints. *Science* 2001;294:2536-9.
- Bouret SG, Draper SJ, Simerly RB. Trophic action of leptin on hypothalamic neurons that regulate feeding. *Science* 2004;304:108-10.

- Brachmann CB, Sherman JM, Devine SE, Carmeron EE, Pillus L, Boeke JD. The SIR2 gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes Dev.* 1995;9:2888-902.
- Brand M, Yamamoto K, Staub A, Tora L. Identification of TATA-binding protein-free TAFII-containing complex subunits suggests a role in nucleosome acetylation and signal transduction. *J. Biol. Chem.* 1999;274:18285-9.
- Branscombe TL, Frankel A, Lee JH, Cook JR, Yang Z, Pestka S, Clarke S. PRMT5 (janu kinase-binding protein 1) catalyzes the formation of symmetric dimethylarginine residues in proteins. *J. Biol. Chem.* 2001;276:32971-6.
- Braun SA, Pantzeter PL, Collinge MA, Althaus FR. Endoglycosidic cleavage of branched polymers by poly(ADP-ribose) glycohydrolase. *Eur. J. Biochem.* 1994;220:369-75.
- Brehm A, Miska A, Reid J, Bannister A, Kouzarides T. The cell cycle-regulating transcription factors E2F-RB. *Br. J. Cancer* 1999;80(S11):38-41.
- Brehm AB, Miska EA, McCance DJ, Reid JL, Bannister AJ, Kouzarides T. Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature* 1998;391:597-601.
- Brenner C. Catalysis in the nitrilase superfamily. *Curr. Opin. Struct. Biol.* 2002;12:775-82.
- Briggs SD, Xiao T, Sun ZW, Caldwell JA, Shabanowitz J, Hunt DF, Allis CD, Strahl BD. Gene silencing: trans-histone regulatory pathway in chromatin. *Nature* 2002;418:498.
- Brochu G, Duchaine C, Thibeault L, Lagueux J, Shah GM, Poirier GG. Mode of action of poly(ADP-ribose) glycohydrolase. *Biochim.Biophys. Acta* 1994;1219:342-50.
- Brownell JE, Zhou J, Ranalli T, Kobayashi R, Edmondson DG, Roth SY, Allis CD. Tetrahymena histone acetyltransferase A: A homolog to yeast Gcn5p linking histone acetylation to gene activation. *Cell* 1996;84:843-51.
- Brueckner B, Boy RG, Siedlecki P, Musch T, Kliem HC, Zielenkiewicz P, Suhai S, Wiessler M, Lyko F. Epigenetic reactivation of tumor suppressor genes by a novel small-molecule inhibitor of human DNA methyltransferases. *Cancer Res.* 2005;65:6305-11.
- Brunet A, Sweeney LB, Sturgill JF, Sturgill JF, Chua KF, Greer PL, Lin Y, Tran H, Ross SE, Mostoslavsky R, Cohen HY, Hu LS, Cheng HL, Jedrychowski MP, Gygi SP, Sinclair DA, Alt FW, Greenberg ME. Stress-dependent regulation of FOXO transcription factors by the SIRT1 deacetylase. *Science* 2004;303:2011-5.
- Burgers WA, Fuks F, Kouzarides T. DNA methyltransferases get connected to chromatin. *Trends Genet.* 2002;18:275-7.
- Cabo SF de, Santos J, Fernandez-Piqueras J. Molecular and cytological evidence of S-adenosyl-L-homocysteine as an innocuous undermethylating agent in vivo. Molecular and cytological evidence of S-adenosyl-L-homocysteine as an innocuous undermethylating agent in vivo. *Cytogenet. Cell Genet.* 1995;71:187-92.
- Camphausen K, Burgan W, Cerra M, Oswald KA, Trepel JB, Lee MJ, Tofilon PJ. Enhanced radiation-induced cell killing and prolongation of gammaH2AX foci expression by the histone deacetylase inhibitor MS-275. *Cancer Res.* 2004;64:316-21.
- Camporeale G, Shubert EE, Sarah G, Cerny R, Zempleni J. K8 and K12 are biotinylated in human histone H4. *Eur. J. Biochem.* 2004;271:2257-63.
- Camporeale G, Suzuki Y, Eissenberg JC, Zempleni J. Low abundance of wild-type biotinidase and holocarboxylase synthetase decreases biotinylation of histones. *FASEB J.* 2005;19:A55.
- Camporeale G, Eissenberg JC, Giordano E, Zempleni J. Lifespan and resistance to heat stress depend on histone biotinylation in *Drosophila melanogaster*. *FASEB J.* 2006;20:A610.
- Cao R, Wang L, Wang H, Xia L, Erdjument-Bromage H, Tempst P, Jones RS, Zhang Y. Role of histone H3 lysine 27 methylation in polycomb-group silencing. *Science* 2002;298:1039-43.
- Cao R, Tsukada Y, Zhang Y. Role of Bmi-1 and Ring1A in H2A ubiquitylation and Hox gene silencing. *Mol. Cell* 2005;20:845-54.

- Cardoso MC, Leonhardt H. DNA methyltransferases is actively retained in cytoplasm during early development. *J. Cell Biol.* 1999;147:25-32.
- Carrozza MJ, Utley RT, Workman JL, Cote J. The diverse functions of histone acetyltransferase complexes. *Trends Genet.* 2003;19:321-9.
- Carruthers LM, Hansen JC. The core histone N termini function independently of linker histones during chromatin condensation. *J. Biol. Chem.* 2000;275:57285-90.
- Cate B ten, Samplonius DF, Bijma T, de Leij LF, Helfrich W, Bremer E. The histone deacetylase inhibitor valproic acid potently augments gemtuzumab ozogamicin-induced apoptosis in acute myeloid leukemia cells. *Leukemia* 2007;21:248-52.
- Cavalli G, Paro R. The Drosophila Fab-7 chromosomal element conveys epigenetic inheritance during mitosis and meiosis. *Cell* 1998;93:505-18.
- Champagne N, Bertos NR, Pelletier N, Wang AH, Vezmar M, Yang Y, Heng HH, Yang XJ. Identification of a human histone acetyltransferase related to monocyte leucemia zinc finger protein. *J. Biol. Chem.* 1999;274:28528-36.
- Chan AO, Boraddus RR, Houlihan PS, Issa JO, Hamilton SR, Rashid A. CpG island methylation in aberrant crypt foci of the colorectum. *Am. J. Pathol.* 2002;160:1823-30.
- Chan SW, Hong W. Retinoblastoma-binding protein 2 (Rbp2) potentiates nuclear hormone receptor-mediated transcription. *J. Biol. Chem.* 2001;276:28402-12.
- Chen D, Ma H, Hong H, Koh SS, Huang SM, Schurter BT, Aswad DW, Stallcup MR. Regulation of transcription by a protein methyltransferase. *Science* 1999;284:2174-7.
- Chen HY, Sun JM, Zhang Y, Davie JR, Meistrich ML. Ubiquitination of histone H3 in elongating spermatids of rat testes. *J. Biol. Chem.* 1998a;273:13165-9.
- Chen RZ, Pettersson U, Beard C, Jackson-Grusby L, Jaenisch R. DNA hypomethylation leads to elevated mutation rates. *Nature* 1998b;395:89-93.
- Chen Z, Clark S, Birkeland M, Sung CM, Lago A, Liu R, Kirkpatrick R, Johanson K, Winkler JD, Hu E. Induction and superinduction of growth arrest and DNA damage gene 45 (GADD45) alpha and beta messenger RNAs by histone deacetylase inhibitors trichostatin A (TSA) and butyrate in SW620 human colon carcinoma cells. *Cancer Lett.* 2002;188:127-40.
- Cheng HL, Mostoslavsky R, Saito S, Manis JP, Gu Y, Patel P, Bronson R, Appella E, Alt FW, Chua KF. Developmental defects and p53 hyperacetylation in Sir2 homolog (SIRT1)-deficient mice. *Proc. Natl. Acad. Sci.* 2003;100:10794-9.
- Cheng JC, Weisenberger DJ, Gonzales FA, Liang G, Xu GL, Hu YG, Marquez VE, Jones PA. Continuous zebularine treatment effectively sustains demethylation in human bladder cancer cells. *Mol. Cell Biol.* 2004a; 24:1270-8.
- Cheng JC, Yoo CB, Weisenberger DJ, Chuang J, Wozniak C, Liang G, Marquez VE, Greer S, Orntoft TF, Thykjaer T, Jones PA. Preferential response of cancer cells to zebulamine. *Cancer Cell* 2004b;6:151-8.
- Cheng X, Collins RE, Zhang X. Structural and sequence motifs to protein (histone) methylation enzymes. *Annu. Rev. Biophys. Biomol. Struct.* 2005;34:267-94.
- Cheung P, Allis CD, Sassone-Corsi P. Signaling to chromatin through histone modifications. *Cell* 2000a; 103:263-71.
- Cheung P, Tanner KG, Cheung WL, Sassone-Corsi P, Denu JM, Allis CD. Synergistic coupling of histone H3 phosphorylation and acetylation in response to epidermal growth factor stimulation. *Mol. Cell* 2000b;5:905-15.
- Chevillard-Briet M, Trouche D, Vandel L. Control of CBP co-activating activity by arginina methylation. *EMBO J.* 2002;21:5457-66.
- Chew YC Raza AS, Sarath G, Zempleni J. Biotinylation of K8 and K12 co-occurs with acetylation and mono-methylation in human histone H4. *FASEB J.* 2006a;20:610.

- Chew YC, Camporeale G, Kothapalli N, Sarath G, Zempleni J. Lysine residues in N- and C-terminal regions of human histone H2A are targets for biotinylation by biotinidase. *J. Nutr. Biochem.* 2006b; 17:225-33.
- Chew YC, Sarath G, Zempleni J. An avidin-based assay for histone debiotinylase activity in human cell nuclei. *J. Nutr. Biochem.* 2007;18:475-81
- Choi SW, Mason JB. Folate status: effects on pathways of colorectal carcinogenesis. *J. Nutr.* 2002;132:2413S-2418S.
- Choi SW, Stickel F, Baik HW, Kim YI, Seitz HK, Mason JB. Chronic alcohol consumption induces genomic but not p53-specific DNA hypomethylation in rat colon. *J. Nutr.* 1999;129:1945-50.
- Christianson AL, Chesler N, Kromberg JG. Fetal valproate syndrome: clinical and neuro-developmental features in two siblings pairs. *Dev. Med. Child Neurol.* 1994;36:361-9.
- CIAQ Inbreeding, genetic principles. *Centre d'insémination artificielle du Québec (CIAQ)*. Disponible en: [http://www.ciaq.com/Consanguinité\\_e.html](http://www.ciaq.com/Consanguinité_e.html). Accesado el 1 de junio 2008.
- Clark SJ, Harrison J, Paul CL, Frommer M. High sensitivity mapping of methylated cytosines. *Nucleic Acids Res.* 1994;22:2990-7.
- Cleary JD, Pearson CE. The contribution od cis-elements to disease-associated repeat instability: clinical and experimental evidence. *Cytogenet. Genome Res.* 2003;100:25-55.
- Clements A, Poux AN, Lo WS, Pillus L, Berger SL, Marmorstein R. Structural basis for histone and phosphohistone binding by the GCN5 histone acetyltransferase. *Mol. Cell* 2003;12:461-73.
- Cohen HY, Miller C, Bitterman KJ, Wall NR, Hekking B, Kessler B, Howitz KT, Gorospe M, de Cabo R, Sinclair DA. Calorie restriction promotes mammalian cell survival by inducing the SIRT1 deacetylase. *Science* 2004;305:390-2.
- Colella S, Shen L, Baggerly KA, Issa JP, Krahe R. Sensitive and quantitative universal pyrosequencing methylation analysis of CpG sites. *Biotechniques* 2003;35:146-50.
- Conaway RC, Brower CS, Conaway JW. Emerging roles of ubiquitina in transcription regulation. *Science* 2002;296:1254-8.
- Conley BA, Wright JJ, Kummar S. Targeting epigenetic abnormalities with histone deacetylase inhibitors. *Cancer* 2006;107:832-40.
- Constância M, Hemberger M, Hughes J, Dean W, Ferguson-Smith A, Fundele R, Stewart F, Kelsey G, Fowden A, Sibley C, Reik W. Placental-specific IGF-II is a major modulator of placental and fetal growth. *Nature* 2002;417:945-8.
- Cook EHJ, Courchesne RY, Cox NJ, Lord C, Gonen D, Guter SJ, Lincoln A, Nix K, Haas R, Leventhal BL, Courchesne E. 1998. Linkage-disequilibrium mapping of autistic disorder, with 15q11-13 markers. *Am. J. Hum. Genet.* 1998;62:1077-83.
- Cook EHJ, Lindgren V, Leventhal BL, Courchesne R, Lincoln A, Shulman C, Lord C, Courchesne E. 1997. Autism or atypical autism in maternally but not paternally derived proximal 15q duplication. *Am. J. Hum. Genet.* 1997;69:928-34.
- Cook JR, Lee JH, Yang ZH, Krause CD, Herth N, Hoffmann R, Pestka S. FBXO11/PRMT9, a new protein arginine methyltransferase, symmetrically dimethylates arginine residues. *Biochem. Biophys. Res. Commun.* 2006;342:472-81.
- Cooney CA. Are somatic cells inherently deficient in methylation metabolism? A proposed mechanism for DNA methylation loss, senescence and aging. *Growth Dev. Aging* 1993;57:261-73.
- Cooney CA, Dave AA, Wolff GL. Maternal methyl supplements in mice affect epigenetic variation and DNA methylation of offspring. *J. Nutr.* 2002;132:S2393-S2400.
- Corda D, Di Girolamo M. Mono-ADP-ribosylation: a tool for modulating immune response and cell signalling. *Sci. STKE* 2002;PE53.
- Corda D, Di Girolamo M. Functional aspects of protein mono-ADP-ribosylation. *EMBO J.* 2003;22:1953-8.
- Costello JF, Fruhwald MC, Smiraglia DJ, DJ, Rush LJ, Robertson GP, Gao X, Wright FA, Feramisco JD, Peltomäki P, Lang JC, Schuller DE, Yu L, Bloomfield CD, Caligiuri MA, Yates A, Nishikawa R, Su Huang

- H, Petrelli NJ, Zhang X, O'Dorisio MS, Held WA, Cavenee WK, Plass C. Aberrant CpG-island methylation has nonrandom and tumour-type-specific patterns. *Nat. Genet.* 2000;24:132-8.
- Costello JF, Plass C, Cavenee WK. Restriction landmark genome scanning. *Methods Mol. Biol.* 2002;200:53-70.
- Covic M, Hassa PO, Saccani S, Buerki C, Meier NI, Lombardi C, Imhof R, Bedford MT, Natoli G, Hottiger MO. Arginine methyltransferase CARM1 is a promoter-specific regulator of NF-kappaB-dependent gene expression. *EMBO J.* 2004;24:85-96.
- Cremer T, Cremer C. Chromosome territories, nuclear architecture and gene regulation in mammalian cells. *Nat. Rev. Genet.* 2001;2:292-301.
- Crisp SERH, Camporeale G, White BR, Tooms CF, Camporeale G, Said HM, Zempleni J. Biotin supply affects rates of cell proliferation, biotinylation of carboxylases and histones, and expression of the gene encoding the sodium-dependent multivitamin transporter in Jar choriocarcinoma cells. *Eur. J. Nutr.* 2004;43:23-31.
- Cummings JH, Pomare EW, Branch WJ, Naylor CP, MacFarlane GT. Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut* 1987;28:1221-7.
- Curley JP, Barton S, Surani A, Keverne EB. Coadaptation in mother and infant regulated by a paternally expressed imprinted gene. *Proc. Biol. Sci.* 2004;271:1303-9.
- Cuthbert GL, Daujat S, Snowden AW, Erdjument-Bromage H, Hagiwara T, Yamada M, Schneider R, Gregory PD, Tempst P, Bannister AJ, Kouzarides T. Histone deimination antagonizes arginine methylation. *Cell* 2004;118:545-53.
- Czermin B, Melfi R, McCabe D, Seitz V, Imhof A, Pirrotta V. Drosophila enhancer Zeste/ESC complexes have a histone H3 methyltransferase activity that marks chromosomal polycomb sites. *Cell* 2002;111:185-96.
- D'Amours D, Desnoyers S, D'Silva I, Poirier GG. Poly(ADP-ribosyl)ation reactions in the regulation of nuclear functions. *Biochem. J.* 1999;342:249-68.
- D'Andrea A, Pellman D. Deubiquitinating enzymes: A new class of biological regulators. *Crit. Rev. Biochem. Mol. Biol.* 1988;33:337-52.
- Dabelea D, Hanson RL, Lindsay RS, Pettitt DJ, Imperatore G, Gabir MM, Roumain J, Bennett PH, Knowler WC. Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. *Diabetes* 2000;49:2208-11.
- Dal Maso L, La Vecchia C, Polesel J, Talamini R, Levi F, Conti E, Zambon P, Negri E, Franceschi S. Alcohol drinking outside meals and cancers to the upper aero-digestive tract. *Int. J. Cancer* 2002;102:435-7.
- Danial NN, Korsmeyer SJ. Cell death: critical control points. *Cell* 2003;116:205-19.
- Daniel H, Wenzel U. Nutritional genomics: concepts, tools and expectations. En: Brigelius-Flohé R, Joost HG (eds). *Nutritional genomics: impact on health and disease*. Alemania: Wiley-VCH; 2006.
- Daniel JA, Torok MS, Sun ZW, Schieltz D, Allis CD, Yates JR, Grant PA. Deubiquitination of histone H2B by a yeast acetyl-transferase complex regulates transcription. *J. Biol. Chem.* 2004;279:1867-71.
- Das PM, Singal R. DNA methylation and cancer. *J. Clin. Oncol.* 2004;22:4632-42.
- Dashwood RH, Myzak M, Ho E. Dietary HDAC inhibitors: time to rethink weak ligands in cancer chemoprevention. *Carcinogenesis* 2006;27:344-9.
- Davie JR, Murphy LC. Level of ubiquitinated histone H2B in chromatin is coupled to ongoing transcription. *Biochemistry* 1999;29:4752-7.
- Davies N, Lindsey GG. Histone H2B (and H2A) ubiquitination allows normal histone octamer and core particle reconstitution. *Biochim. Biophys. Acta* 1994; 1218:187-93.
- Davis AJ, Gelmon KA, SiuLL, Moore MJ, Britten CD, Mistry N, Klamut H, D'Aloisio S, MacLean M, Wainman N, Ayers D, Firby P, Besterman JM, Reid GK, Eisenhauer EA. Phase I and pharmacologic study of the human DNA methyltransferase antisense oligodeoxynucleotide MG98 given as a 21-day continuous infusion every 4 weeks. *Invest. New Drugs* 2003a; 21:85-7.
- Davis CA, Haberland M, Arnold MA, Sutherland, LB, McDonald OG, Richardson JA, Childs G, Harris S, Owens GK, Olson EN. PRISM/PRDM6, a transcriptional repressor that promotes the proliferative gene program in smooth muscle cells. *Mol. Cell Biol.* 2006; 26:2626-36.

- Davis CD, Uthus EO. Dietary folate and selenium affect dimethylhydrazine-induced aberrant crypt formation, global DNA methylation and one-carbon metabolism in rats. *Nutr.* 2003; 133:2907-14.
- Davis CD, Uthus EO. DNA methylation, cancer susceptibility, and nutrient interactions. *Exp. Biol. Med. (Maywood)* 2004; 229:988-95.
- Davis DC, Uthus EO, Finley JW. Dietary selenium and arsenic affect DNA methylation in vitro in Caco-2 cells and in vivo in rat liver and colon. *J. Nutr.* 2000; 130:2903-9.
- Davuluri RV, Grosse I, Zhang MQ. Computational identification of promoters and first exons in the human genome. *Nat. Genet.* 2001;29:412-7.
- Dawson BA, Herman T, Haas AL, Lought J. Affinity isolation of active murine erythroleukemia cell chromatin: Uniform distribution of ubiquitinated histone H2A between active and inactive fractions. *J. Cell Biochem.* 1991;46:166-73.
- Day JK, Bauer AM, desBordes C, Zhuang Y, Kim BE, Newton LG, Nehra V, Forsee KM, MacDonald RS, Besch-Williford C, Huang TH, Lubahn DB. Genistein alters methylation patterns in mice. *J. Nutr.* 2002;132:2419S-2432S.
- Delaval K, Feil R. Epigenetic regulation of mammalian genomic imprinting. *Curr. Opin. Genet.Dev.* 2004;14:188-93.
- Delaval K, Wagschal A, Feil R. Epigenetic deregulation of imprinting in congenital diseases of aberrant growth. *Bioessays* 2006; 28:453-9.
- Delrue MA, Michaud JL. Fat chance: genetic syndromes with obesity. *Clin. Genet.* 2004;66:88-93.
- Demenais F, Chaudru V, Marínez M. Detection of parent-of-origin effects for atopy by model-free and model-based linkage analyses. *Genet. Epigemiol.* 2001;21:S186-S191.
- Deng C, Kaplan MJ, Yang J, Ray D, Zhang Z, McCune WJ, Hanash SM, Richardson BC. Decreased Ras-mitogen-activated protein kinase signaling may cause DNA hypomethylation in T lymphocytes from lupus patients. *Arthritis Rheum.* 2001;44:397-407.
- Deng C, Lu Q, Zhang Z, Rao T, Attwood J, Yung R, Richardson B. Hydralazine may induce autoimmunity by inhibiting extracellular signal-regulated kinase pathway signaling. *Arthritis Rheum.* 2003;48:746-56.
- Deng Q, Huang S. PRDM5 is silenced in human cancers and has growth suppressive activities. *Oncogene* 2004;23:4903-10.
- Denu JM. The Sir2 family of protein deacetylases. *Curr. Opin. Chem. Biol.* 2005;9:431-40.
- Derunes C, Briknarova K, Geng L, Lib S, Gessnerb CR, Hewittb K, Wu SD, Huanga S, Woods VI Jr., Ely KR. Characterization of the PR domain of RIZ1 histone methyltransferase. *Biochem. Biophys. Res. Commun.* 2005;333:925-34.
- Desmarais Y, Menard L, Lagueux J, Poirier GG. Enzymological properties of poly(ADP-ribose)polymerase : characterization of automodification sites and NADase activity. *Biochim. Biophys. Acta* 1991;1078:179-86.
- Di Croce L, Raker VA, Corsaro M, Fazi F, Fanelli M, Faretta M, Fuks F, Lo Coco F, Kouzarides T, Nervi C, Minucci S, Pelicci PG. Methyltransferase recruitment and DNA hypermethylation of target promoters by an oncogenic transcription factor. *Science* 2002;295:1079-82.
- Di Fiore PP, Polo S, Hofmann K. When ubiquitin meets ubiquitin receptors: A signalling connection. *Nat. Rev. Mol. Cell Biol.* 2003;4:491-7.
- Di Meglio S, Denegri M, Vallefuoco S, Tramontano F, Scovassi AI, Quesada P. Poly(ADPR) polymerase-1 and poly(ADPR) glycohydrolase level and distribution in differentiating rat germinal cells. *Mol. Cell Biochem.* 2003;248:85-91.
- Doherty AS, Bartolomei MS, Schultz RM. Regulation of stage-specific nuclear translocation of Dnmt1o during preimplantation mouse development. *Dev. Biol.* 2002;242:255-66.
- Dokmanovic M, Marks PA. Prospects: histone deacetylase inhibitors. *J. Cell Biochem.* 2005;96:293-304.
- Dokmanovic M, Clarke C, Marks PA. Histone deacetylase inhibitors: overview and perspectives. *Mol. Cancer Res.* 2007;5:981-9.

- Domínguez G, Carballido J, Silva J, Silva JM, García JM, Menéndez J, Provencio M, España P, Bonilla F. P14ARF promoter hypermethylation in plasma DNA as an indicator of disease recurrence in bladder cancer patients. *Clin. Cancer Res.* 2002;8:980-5.
- Dong A, Yoder JA, Zhang X, Zhou L, Bestor TH, Cheng X. Structure of human DNMT2, an enigmatic DNA methyltransferase homolog that displays denaturant-resisting binding to DNA. *Nucleic Acids Res.* 2001;29:439-48.
- Dong C, Yoon W, Goldschmidt-Clermont PJ. DNA methylation and atherosclerosis. *J. Nutr.* 2002;132:2406S-2409S.
- Dou Y, Milne TA, Tackett AJ, Smith ER, Fukuda A, Wysocka J, Allis CD, Chait BT, Hess JL, Roeder RG. Physical association and coordinate function of the H3 K4 methyltransferase MLL1 and the H4 K16 acetyltransferase MOF. *Cell* 2005;121:873-85.
- Dover J, Schneider J, Tawiah-Boateng MA, Wood A, Dean K, Johnston M, Shilatifard A. Methylation of histone H3 by COMPASS requires ubiquitination of histone H2B by Rad6. *J. Biol. Chem.* 2002;277:28368-71.
- Downs JA, Lowndes NF, Jackson SP. A role for *Saccharomyces cerevisiae* histone H2A in DNA repair. *Nature* 2000;408:1001-4.
- Duesne N, Pagniez A, Mayeur C, Thomas M, Cherbuy C, Duée PH, Martel P, Chaumontet C. Diallyl disulfide (DADS) increases histone acetylation and p21<sup>Waf1/Cip1</sup> expression in human colon tumor cell lines. *Carcinogenesis* 2004;25:1227-36.
- Duggan DJ, Bittner M, Chen Y, Meltzer P, Trent JM. Expression profiling using cDNA microarrays. *Nature Genet.* 1999;21:10-4.
- Duvic M, Talpur R, Ni X, Zhang C, Hazarika P, Kelly C, Chiao JH, Reilly JF, Ricker JL, Richon VM, Frankel SR. Phase 2 trial of oral vorinostat (suberoylanilide hydroxamic acid, SAHA) for refractory cutaneous T-cell lymphoma (CTCL). *Blood* 2007;109:31-9.
- Dyson N. The regulation of E2F by pRB-family proteins. *Genes Dev.* 1998;12:2245-62.
- Eads CA, Danenberg KD, Kawakami K, Saltz LB, Blake C, Shibata D, Danenberg PV, Laird PW. MethylLight: a high-throughput assay to measure DNA methylation. *Nucleic Acids Res.* 2000;28:E32.
- Edberg DD, Bruce JE, Siems WF, Reeves R. In vivo posttranslational modifications of the high mobility group A1A proteins in breast cancer cells of differing metastatic potential. *Biochemistry* 2004;43:11500-15.
- Eden A, Gaudet F, Waghmare A, Jaenisch R. Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 2003;300:455.
- Egger G, Liang G, Aparicio A, Jones PA. Epigenetics in human disease and prospects for epigenetic therapy. *Nature* 2004;429:457-63.
- Ehrlich M. DNA methylation in cancer: too much, but also too little. *Oncogene* 2002;21:5400-13.
- Ehrlich M, Gama-Sosa MA, Huang LH, Midgett RM, Kuo KC, McCune RA, Gehrke C. Amount and distribution of 5-methylcytosine in human DNA from different types of tissues of cells. *Nucleic Acids Res.* 1982;10:2709-21.
- Ehrlich M, Nelson MR, Stanssens P, Zabeau M, Liloglou T, Xinarianos G, Cantor CR, Field JK, van den Boom D. Quantitative high-throughput analysis of DNA methylation patterns by base-specific cleavage and mass spectrometry. *Proc. Natl. Acad. Sci. USA* 2005;102:15785-90.
- Engeland M van, Weijenberg MP, Roemen GM, Brink M, Bruïne AP de, Goldbohm RA, Brandt PA van den, Baylin SB, Goeij AFPM de, Herman JG. Effects of dietary folate and alcohol intake on promoter methylation in sporadic colorectal cancer: the Netherlands cohort study on diet and cancer. *Cancer Res.* 2003;63:3133-7.
- Ergul E, Sazci A, Utkan Z, Canturk NZ. Polymorphisms in the MTHFR gene are associated with breast cancer. *Tumor Biol.* 2003;24:286-90.
- Ericsson C, Goldknopf IL, Daneholt B. Inhibition of transcription does not affect the total amount of ubiquitinated histone 2A in chromatin. *Exp. Cell. Res.* 1986;167:127-34.
- Ernst P, Wang J, Huang M, Goodman RH, Korsmeyer SJ. MLL and CREB bind cooperatively to the nuclear coactivator CREB-binding protein. *Mol. Cell Bio.* 2001;21:2249-58.

- Esteller M. DNA methylation and cancer therapy: new developments and expectations. *Curr. Opin. Oncol.* 2005;17:55-60.
- Esteller M. Cancer epigenomics: DNA methylomes and histone modification maps. *Nat. Rev. Genet.* 2007;8:286-98.
- Esteller M. Epigenetics in cancer. *N. Engl. J. Med.* 2008;358:1148-59.
- Esteller M, Herman JG. Generating mutations but providing chemosensitivity: the role of O6-methylguanine DNA methyltransferase in human cancer. *Oncogene* 2004;23:1-8.
- Esteller M, Sánchez-Céspedes M, Rosell R, Sidransky DB, Baylin SB, Herman JG. Detection of aberrant promoter hypermethylation of tumor suppressor genes in serum DNA from non-small cell lung cancer patients. *Cancer Res.* 1999a;59:67-70.
- Esteller M, Hamilton SR, Burger PC, Baylin SB, Herman JG. Inactivation of the DNA repair gene O6-methylguanine-DNA methyltransferase by promoter hypermethylation is a common event in primary human neoplasia. *Cancer Res.* 1999b;59:793-7.
- Esteller M, García-Foncillas J, Andion E, Goodman SN, Hidalgo OF, Vanaclocha V, Baylin SB, Herman JG. Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents. *N. Engl. J. Med.* 2000; 343:1350-4.
- Esteller M, Corn PG, Baylin SB, Herman JG. A gene hypermethylation profile of human cancer. *Cancer Res.* 2001a;61:3225-9.
- Esteller M, Fraga MF, Guo M, Garcia-Foncillas J, Hedenfalk I, Godwin AK, Trojan J, Vaurs-Barrière C, Bignon YJ, Ramus S, Benitez J, Caldes T, Akiyama Y, Yuasa Y, Launonen V, Canal MJ, Rodriguez R, Capella G, Peinado MA, Borg A, Aaltonen LA, Ponder BA, Baylin SB, Herman JG. DNA methylation patterns in hereditary human cancers mimic sporadic tumorigenesis. *Hum. Mol. Genet.* 2001b;10:3001-7.
- Esteller M, Gaidano G, Goodman SN, Zagonel V, Capello D, Botto B, Rossi D, Gloghini A, Vitolo U, Carbone A, Baylin SB, Herman JG. Hypermethylation of the DNA repair gene O(6)-methylguanine DNA methyltransferase and survival of patients with diffuse large B-cell lymphoma. *J. Natl. Cancer Inst.* 2002;94:26-32.
- Evron E, Dooley WC, Umbricht CB, Rosenthal D, Sacchi N, Gabrielson E, Soito AB, Hung DT, Ljung B, Davidson NE, Sukumar S. Detection of breast cancer cells in ductal lavage fluid by methylation-specific PCR. *Lancet* 2001;357:1335-6.
- Fang JY, Xiao SD. Folic acid, polymorphism of methyl-group metabolism genes, and DNA methylation in relation to GI carcinogenesis. *Gastroenterol.* 2003;38:821-9.
- Fang MZ, Wang Y, Ai N, Hou Z, Sun Y, Lu H, Welsh W, Yang CS. Tea polyphenol (-)-epigallocatechin-3-gallate inhibits DNA methyltransferase and reactivates methylation-silenced genes in cancer cell lines. *Cancer Res.* 2003;63:7563-70.
- Fatemi SH, Earle JA, McMenomy T. Reduction in Reelin immunoreactivity in hippocampus of subjects with schizophrenia, bipolar disorder and major depression. *Mol. Psychiatry* 2000;5:654-63.
- Fatemi SH, Snow AV, Stary JM, Araghi-Niknam M, Reutiman TJ, Lee S, Brooks AI, Pearce DA. Reelin signaling is impaired in autism. *Biol. Psychiatry* 2005;57:777-87.
- Fears S, Mathieu C, Zelezniak-Le N, Huang S, Rowley JD, Nucifora G. Intergenic splicing of MDS1 and EVI1 occurs in normal tissue as well as in myeloid leukaemia and produces a new member of the PR domain family. *Proc. Natl. Acad. Sci. USA* 1996;93:1642-7.
- Feil R. Environmental and nutritional effects on the epigenetic regulation of genes. *Mutat. Res.* 2006; 600:46-57.
- Feinberg AP. Epigenetics at the epicenter of modern medicine. *JAMA* 2008; 299:1345-50.
- Feinberg AP. Phenotypic plasticity and the epigenetics of human disease. *Nature* 2007; 447:433-40.
- Feinberg AP, Tycko B. The history of cancer epigenetics. *Nat. Rev. Cancer* 2004;4:143-53.
- Feinberg AP, Cul H, Ohlsson R. DNA methylation and genomic imprinting: insights from cancer into epigenetic mechanisms. *Sem. Canc. Biol.* 2002;12:389-98.
- Feinberg AP, Ohlsson R, Henikoff S. The epigenetic progenitor origin of human cancer. *Nat. Rev. Genet.* 2006;7:21-3.

- Feng Q, Zhang Y. The MeCP1 complex represses transcription through preferential binding, remodeling, and deacetylating methylated nucleosomes. *Genes Dev.* 2001;15:827-32.
- Feng Q, Wang H, Ng HH, Erdjument-Bromage H, Tempst P, Struhl K, Zhang Y. Methylation of H3-lysine 79 is mediated by a new family of HMTases without a SET domain. *Curr. Biol.* 2002;12:1052-8.
- Ferreri AJ, Dell'Oro S, Capello D, Ponzone M, Iuzzolino P, Rossi D, Pasini F, Ambrosetti A, Orvieto E, Ferrarese F, Arrigoni G, Foppoli M, Reni M, Gaidano G. Aberrant methylation in the promoter region of the reduced folate carrier gene is a potential mechanism of resistance to methotrexate in primary central nervous system lymphomas. *Br. J. Haematol.* 2004;126:657-64.
- Ferro AM, Olivera BM. Poly(ADP-ribosylation) in vitro. Reaction parameters and enzyme mechanism. *J. Biol. Chem.* 1982;257:7808-13.
- Finnin MS, Dongian JR, Cohen A, Richon VM, Rifkind RA, Marks PA, Breslow R, Pavletich NP. Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors. *Nature* 1999;401:188-93.
- Fischer DD, Cai R, Bhatia U, Asselbergs FA, Song C, Terry R, Trogani N, Widmer R, Atadja P, Cohen D. Isolation and characterization of a novel class II histone deacetylase, HDAC10. *J. Biol. Chem.* 2002;277:6656-66.
- Fischle W, Dequiedt F, Fillion M, Hendzel MJ, Voelter W, Verdin E. Human HDAC7 histone deacetylase activity is associated with HDAC3 in vivo. *J. Biol. Chem.* 2001;276:35826-35.
- Fischle W, Dequiedt F, Hendzel MJ, Guenther MG, Lazar MA, Voelter W, Verdin E. Enzymatic activity associated with class II HDACs is dependent on a multiprotein complex containing HDAC3 and SMRT/N-CoR. *Mol. Cell* 2002;9:45-57.
- Fischle W, Wang Y, Allis CD. Histone and chromatin cross-talk. *Curr. Opin. Cell Biol.* 2003a; 15:172-83.
- Fischle W, Wang Y, Jacobs SA, Kim Y, Allis CD, Khorasanizadeh S. Molecular basis for the discrimination of repressive methyl-lysine marks in histone H3 by polycomb and HP1 chromodomains. *Genes Dev.* 2003b;17:1870-81.
- Fodor BD, Kubicek S, Yonezawa M, O'Sullivan RJ, Sengupta R, Perez-Burgos L, Opravil S, Mechteder K, Schotta G, Jenuwein T. Jmjd2b antagonizes H3K9 trimethylation at pericentric heterochromatin in mammalian cells. *Genes Dev.* 2006;20:1557-62.
- Ford E, Voit R, Liszt G, Magin C, Grummt I, Guarente L. Mammalian Sir2 homolog SIRT7 is an activator of RNA polymerase I transcription. *Genes Dev.* 2006;20:1075-80.
- Fouladi M, Furman WL, Chin T, Freeman BB 3rd, Dudkin L, Stewart CF, Krailo MD, Speights R, Ingle AM, Houghton PJ, Wright J, Adamson PC, Blaney SM. Phase I study of depsipeptide in pediatric patients with refractory solid tumors: a Children's Oncology Group report. *J. Clin. Oncol.* 2006;24:3678-85.
- Fraga MF, Ballestar E, Paz MF, Ropero S, Setien F, Ballestar ML, Heine-Suñer D, Cigudosa JC, Urioste M, Benitez J, Boix-Chornet M, Sanchez-Aguilera A, Ling C, Carlsson E, Poulsen P, Vaag A, Stephan Z, Spector TD, Wu YZ, Plass C, Esteller M. Epigenetic differences arise during the lifetime of monozygotic twins. *Proc. Natl. Acad. Sci. USA* 2005a;102:10604-9.
- Fraga MF, Ballestar E, Villar-Garea A, Boix-Chornet M, Espada J, Schotta G, Bonaldi T, Haydon C, Ropero S, Petrie K, Iyer NG, Pérez-Rosado A, Calvo E, Lopez JA, Cano A, Calasanz MJ, Colomer D, Piris MA, Ahn N, Imhof A, Caldas C, Jenuwein T, Esteller M. Loss of acetylation at Lys16 and trimethylation at Lys20 of histone H4 is a common hallmark of human cancer. *Nat. Genet.* 2005;37:391-400.
- Frechette A, Huletsky A, Aubin RJ, de Murcia G, Mandel P, Lord A, Grondin G, Poirier GG. Poly(ADP-ribosylation) of chromatin: kinetics of relaxation and its effect on chromatin solubility. *Can. J. Biochem. Cell Biol.* 1985;63:764-73.
- Freeman-Cook LL, Gomez EB, Spedale EJ, Marlett J, Forsburg SL, Pillus L, Laurenson P. Conserved locus-specific silencing functions of *Schizosaccharomyces pombe* Sir2+. *Genetics* 2005;169:1243-60.
- Freudenheim JL, Bonner M, Krishnan S, Ambrosone CB, Graham S, McCann SE, Moysich KB, Bowman E, Nemoto T, Shields PG. Diet and alcohol consumption in relation to p53 mutations in breast tumors. *Carcinogenesis* 2004;25:931-9.

- Frigola J, Song J, Stirzaker C, Hinshelwood RA, Peinado MA, Clark SJ. Epigenetic remodeling in colorectal cancer results in coordinate gene suppression across an entire chromosome band. *Nat. Genet.* 2006;38:540-9.
- Frommer M, McDonald LE, Milar DS, Collis CM, Watt F, Grigg GW, Molloy PL, Paul CL. A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands. *Proc. Natl. Acad. Sci. USA* 1992;89:1827-31.
- Frye RA. Characterization of five human cDNAs with homology to yeast SIR2 gene: Sir-2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity. *Biochem. Biophys. Res. Commun.* 1999;260:273-9.
- Frye RA. Phylogenetic classification of prokaryotic and eukaryotic Sir2-like proteins. *Biochem. Biophys. Res. Commun.* 2000;273:793-8.
- Fuchs CS, Willet WC, Colditz GA, Hunter DJ, Stampfer MJ, Speizer FE, Giovannucci EL. The influence of folate and multivitamin use on the familial risk of colon cancer in women. *Cancer Epidemiol. Biomarkers Prev.* 2002;11:227-34.
- Fuks F, Burgers WA, Brehm A, Hughes-Davies L, Kouzarides T. DNA methyltransferases Dnmt1 associates with histone deacetylase activity. *Nature Genet.* 2000;24:88-9.
- Fuks F, Burgers WA, Godin N, Kasai M, Kouzarides T. DNMT3a binds deacetylases and is recruited by a sequence-specific repressor to silence transcription. *EMBO J.* 2001;20:2536-44.
- Fulco M, Schiltz RL, Iezzi S, King MT, Zhao P, Kashiwaya Y, Hoffman E, Veech RL, Sartorelli V. Sir2 regulates skeletal muscle differentiation as a potential sensor of the redox state. *Mol. Cell* 2003;51-62.
- Furumai R, Komatsu Y, Nishino N, Khochbin S, Yoshida M, Hironouchi S. Potent histone deacetylase inhibitors built from trichostatin A and cyclic tetrapeptide antibiotics including trapoxin. *Proc. Natl. Acad. Sci. USA* 2001;98:87,92.
- Futscher BW, O'Meara MM, Kim CJ, Rennels MA, Lu D, Gruman LM, Seftor REB, Hendrix MJC, Domann FE. Aberrant methylation of the maspin promoter is an early event in human breast cancer. *Neoplasia* 2004;6:380-9.
- Gallou-Kabani C, Junien C. Nutritional epigenomics of metabolic syndrome. New perspective against the epidemic. *Diabetes* 2005;1899-906.
- Gao L, Cueto MA, Asselbergs F, Atadja P. Cloning and functional characterization of HDAC11, a novel member of the human histone deacetylase family. *J. Biol. Chem.* 2002;277:25748-55.
- Garcia-Manero G, Kantarjian HM, Sanchez-Gonzalez B, Yang H, Rosner G, Verstovsek S, Rytting M, Wierda WG, Ravandi F, Koller C, Xiao L, Faderl S, Estrov Z, Cortes J, O'Brien S, Estey E, Bueso-Ramos C, Fiorentino J, Jabbour E, Issa JP. Phase 1/2 study of the combination of 5-aza-2'-deoxycytidine with valproic acid in patients with leucemia. *Blood* 2006;108:3271-9.
- Garcia-Salcedo JA, Gijon P, Nolan DP, Tebabi P, Pays E. A chromosomal SIR2 homologue with both histone NAD-dependent ADP-ribosyltransferase and deacetylase activities is involved in DNA repair in *Tripanosoma brucei*. *EMBO J.* 2003;22:5851-62.
- Gary JD, Clarke S. RNA and protein interactions modulated by protein arginine methylation. *Prog. Nucleic Acids Res. Mol. Biol.* 1998;61:65-131.
- Georgakopoulos T, Thireos G. Two distinct yeast transcriptional activators require the function of the GCN5 protein to promote normal levels of transcription. *EMBO J.* 1992;11:4145-52.
- Gill G. Post-translational modification by the small ubiquitin-related modifier SUMO has big effects on transcription factor activity. *Curr. Opin. Genet. Dev.* 2003; 13:108-113.
- Gill G. SUMO and ubiquitin in the nucleus: different functions, similar mechanisms? *Genes Dev.* 2004;18:2046-59.
- Gleave ME, Monia BP. Antisense therapy for cancer. *Nat. Rev. Cancer* 2005;5:468-79
- Glick BR, Pasternak JJ. *Molecular Biotechnology – Principles and Applications of Recombinant DNA*. Washington: ASM Press; 2005.

- Glowacki G, Braren R, Firner K, Nissen M, Kühl M, Reche P, Bazan F, Cetkovic-Cvrlje M, Leiter E, Haag F, Koch-Nolte F. The family of toxin-related ecto-ADP-ribosyltransferases in humans and the mouse. *Protein Sci.* 2002;11:1657-70.
- Gluckman PD, Hanson MA, Beedle AS. Non-genomic transgenerational inheritance of disease risk. *Bioessays* 2007;29:145-54.
- Goebel MG, Yochem J, Jentsch S, McGrath JP, Varshavsky A, Byers B. The yeast cell cycle gene CDC34 encodes a ubiquitina-conjugating enzyme. *Science* 1988;241:1331-5.
- Goessl C, Krause H, Muller M, Heicappell R, Schrader M, Sachsinger J, Miller K. Fluorescent methylation-specific polymerase chain reaction for DNA-based detection of prostate cancer in bodily fluids. *Cancer Res.* 60;5941-5,
- Goff DC, Bottiglieri T, Arning E, Shih V, Freudenreich O, Evins AE, Henderson DC, Baer L, Coyle J. Folate, homocysteine, and negative symptoms in schizophrenia. *Am. J. Psychiatry* 2004;1705-8.
- Gojo I, Jiemitz A, Trepel JB, Sparreboom A, Figg WD, Rollins S, Tidwell ML, Greer J, Chung EJ, Lee MJ, Gore SD, Sausville EA, Zwiebel J, Karp JE. Phase 1 and pharmacological study of MS-275, a histone deacetylase inhibitor, in adults with refractory and relapsed acute leukemias. *Blood* 2006;109:2781-90.
- Golderer G, Grobner P. ADP-ribosylation of core histones and their acetylated subspecies. *Biochem J* 1991; 277:607-10.
- Golderer G, Loidl P, Grobner P. Cell-cycle-dependent ADP-ribosylation of the nuclear matrix. *Eur. J. Cell. Biol.* 1991; 55:183-5.
- Goldknopf IL, Taylor CW, Baum RM, Yeoman LC, Olson MO, Prestayko AW, Busch H. Isolation and characterization of protein A24, a "histone like" non-histone chromosomal protein. *J. Biol. Chem.* 1975;250:7182-7.
- Goll MG, Bestor TH. Eukaryotic cytosine methyltransferases. *Annu. Rev.* 2005;74:481-514.
- Gong L, Millas S, Maul GG, Yeh ET. Differential regulation of sentrinized proteins by a novel sentrin-specific protease. *J. Biol. Chem.* 2000;275:3355-9.
- Gordon, W. Polymers & Liquid Crystals Main Page. Case Western Reserve University. Disponible en: <http://plc.cwru.edu/tutorial/enhanced/main.htm>. Accesado el 10 de junio 2008.
- Gowher H, Jeltsch A. Enzymatic properties of recombinant Dnmt3a DNA methyltransferase from the mouse: the enzyme modifies DNA in a non-processive manner and also methylates non-CpG sites. *J. Mol. Biol.* 2001;309:1201-8.
- Grant PA. A tale of histone modifications. *Genome Biol.* 2001;2(4):1-6.
- Grant PA, Duggan L, Côté J, Roberts SM, Brownell JE, Candau R, Ohba R, Owen-Hughes T, Allis CD, Winston F, Berger SL, Workman JL. Yeast Gcn5 functions in two multisubunit complexes to acetylate nucleosomal histones: Characterization of an Ada complex and the SAGA (Spt9/Ada) complex. *Genes Dev.* 1997;11:1640-50.
- Grant PA, Schiltz D, Pray-Grant MG, Yates JR, Workman JL. The ATM-related factor Tra1 is a component of the purified SAGA complex. *Mol. Cell* 1998;2:863-7.
- Grant S, Easley C, Kirkpatrick P. Vorinostat. *Nat. Rev. Drug Discov.* 2007;6:21-2.
- Gray SG, Qian CN, Furge K, Guo X, Teh BT. Microarray profiling of the effects of histone deacetylase inhibitors on gene expression in cancer cell lines. *Int. J. Oncol.* 2004;24:773-95.
- Gray SG, Iglesias AH, Lizcano F, Villanueva R, Camelo S, Jingu H, Teh BT, Koibuchi N, Chin WW, Kokkotou E, Dangond F. Functional characterization of JMJD2A, a histone deacetylase- and retinoblastoma-binding protein. *J. Biol. Chem.* 2005;280:28507-15.
- Grayson DR, Jia X, Chen Y, Sharma RP, Mitchell CP, Guidotti A, Costa E. Reelin promoter hypermethylation in schizophrenia. *Proc. Natl. Acad. Sci. USA* 2005;102:9341-6.
- Grayson DR, Chen Y, Costa E, Dong E, Guidotti A, Kundakovic M, Sharma RP. The human reelin gene: transcription factors (+), repressors (-) and the methylation switch (+/-) in schizophrenia. *Pharmacol. Ther.* 2006;111:272-86.

- Grbavec D, Lo R, Liu Y, Greenfield A, Stifani S. Groucho/transducin-like enhancer of split (TLE) family members interact with the yeast transcriptional co-repressor SSN6 and mammalian SSN6-related proteins: implication or evolutionary conservation of transcription repression mechanisms. *Biochem. J.* 1999;337:13-7.
- Griffiths AJF, Gelbart WM, Lewontin RC, Miller JH. *Modern Genetic Analysis: Integrating Genes and Genomes*. WH Freeman; 2002.
- Grozinger CM, Schreiber SL. Regulation of histone deacetylase 4 and 5 and transcripcional activity by 14-3-3-dependent cellular localization. *Proc. Natl. Acad. Sci. USA* 2000;97:7835-40.
- Grubisha O, Smith BC, Denu JM. Small molecule regulation of Sir2 protein deacetylases. *FEBS J.* 2005;272:4607-16.
- Grunstein M. Histone acetylation in chromatin structure and transcription. *Nature* 1997;389:349-52.
- Guenther MG, Jenner RG, Chevalier B, Nakamura T, Croce CM, Canaani E, Young RA. Global and Hox-specific roles for the MLL1 methyltransferase. *Proc. Natl. Acad. Sci. USA* 2005;102:8603-8.
- Guidotti A, Auta J, Davis JM, Di-Giorgi-Gerevini V, Dwivedi Y, Grayson DR, Impagnatiello F, Pandey G, Pesold C, Sharma R, Uzunov D, Costa E. Decrease in reelin and glutamic acid decarboxylase67 (GAD67) expression in schizophrenia and bipolar disorder: a postmortem brain study. *Arch. Gen. Psychiatry* 2000;57:1061-9.
- Guo YF, Shen H, Liu YJ, Wang W, Xiong DH, Xiao P, Liu YZ, Zhao LJ, Recker RR, Deng HW. Assessment of genetic linkage and parent-of-origin effects in obesity. *J. Clin. Endocrinol. Metab.* 2006;91:4001-5.
- Guyonnet D, Berges R, Siess MH, Pinnert MF, Chagnon MC, Suschetet M, Le Bon AM. Post-initiation modulating effects of allyl sulfides in rat hepatocarcinogenesis. *Food Chem. Toxicol.* 2004;42:1479-85.
- Gyory I, Wu J, Fejer G, Seto E, Wright KL. PRDI-BF1 recruits the histone H3 methyltransferase G9a in transcriptional silencing. *Nat. Immunol.* 2004;5:299-308.
- Haas AL, Siepmann TJ. Pathways of ubiquitina conjugation. *FASEB J.* 1997;11:1257-68.
- Haas AL, Warms JV, Herskho A, Rose IA. Ubiquitin-activating enzyme. Mechanism and role in protein-ubiquitin conjugation. *J. Biol. Chem.* 1982;257:2543-8.
- Haas AL, Reback PB, Chau V. Ubiquitin conjugation by the yeast RAD6 and CDC34 gene products. Comparison to their putative rabbit homologs, E2(20K) and E2(32K). *J. Biol. Chem.* 1991;266:5104-12.
- Hadorn E. Transdetermination in cells. *Sci. Am.* 1978;219:110-4.
- Haigis MC, Guarente LP. Mammalian sirtuins – emerging roles in physiology, aging and calorie restriction. *Genes Dev.* 2006;20:2913-21.
- Haince JF, Ouellet ME, McDonald D, Hendzel MJ, Poirier GG. Dynamic relocation of poly(ADP-ribose) glycohydrolase isoforms during radiation-induced DNA damage. *Biochem. Biophys. Acta* 2006; 176:226-37.
- Hakimi MA, Bochar DA, Chenoweth J, Lane WS, Mandel G, Shickhattar R. A core-BRAF35 complex containing histone deacetylase mediates repression of neuronal-specific genes. *Proc. Natl. Acad. Sci. USA* 2002;99:7420-5.
- Hales CN, Barker DJ. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 1992;35:595-601.
- Haliwell B. Vitamin C and genomic stability. *Mutat. Res.* 2001;475:29-35.
- Hall IM, Shankaranarayana GD, Norma K, Ayoub N, Cohen A, Gewall SI. Establishment and maintenance of a heterochromatin domain. *Science* 2002; 297:2232-7.
- Hall JG. Genomic imprinting: review and relevance to human diseases. *Am. J. Hum. Genet.* 1990; 46:857-73.
- Hamamoto R, Furukawa Y, Morita M, Iimura Y, Silva FP, Li M, Yagyu R, Nakamura Y. SMYD3 encodes a histone methyltransferase involved in the proliferation of cancer cells. *Nat. Cell Biol.* 2004;6:731-40.
- Han Z, Gou L, Wang H, Shen Y, Deng XW, Chai J. Structural basis for the specific recognition of methylated histone H3 lysine 4 by the WD-40 protein WDR5. *Mol. Cell* 2006;22:137-144.
- Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000;100:57-70.

- Hang J, Dasso M. Association of the human SUMO-1 protease SENP2 with the nuclear pore. *J. Biol. Chem.* 2002;277:19961-6.
- Hanson R, Hess JS, Yu BD, Ernst P, van Lohuizen M, Berns A, van der Lugt NM, Shashikant CS, Ruddle FH, Seto M, Korsmeyer SJ. Mammalian trithorax and polycomb-group homologues are antagonistic regulators of homeotic development. *Proc. Natl. Acad. Sci. USA* 1999;96:14372-7.
- Hao Y, Crenshaw T, Moulton T, Newcomb E, Tycko B. Tumour-suppressor activity of H19 RNA. *Nature* 1993;365:764-7.
- Harmon CE, Portanova JP. Drug-induced lupus: clinical and serological studies. *Clin. Rheum. Dis.* 1982;8:121-35.
- Harms KL, Chen X. Histone deacetylase 2 modulates p53 transcriptional activities through regulation of p53-DNA binding activity. *Cancer Res.* 2007;67:3145-52.
- Harnack L, Jacobs DRJ, Nicodemus K, Lazovich D, Anderson K, Folsom AR. Relationship of folate, vitamin B-6, vitamin B-12, and methionine intake to incidence of colorectal cancers. *Nutr. Cancer* 2002;43:152-8.
- Hassa PO, Hottiger MO. The functional role of poly(ADP-ribose)polymerase 1 as a novel coactivator of NF-kappaB in inflammatory disorders. *Cell Mol. Life Sci.* 2002;59:1534-53.
- Hassa PO, Hottiger MO. An epigenetic code for DNA damage repair pathways? *Biochem. Cell Biol.* 2005;85:270-85.
- Hassa PO, Haenni SS, Elser M, Hottiger MO. Nuclear ADP-ribosylation reactions in mammalian cells: where are we today and where are we going? *Microbiol. Mol. Biol. Rev.* 2006;70:789-829.
- Hatchwell E, Greally JM. The potential role of epigenomic dysregulation in complex human disease. *Trends Genet.* 2007; 23:588-95.
- Hau P, Stupp R, Hegi ME. MGMT methylation status: the advent of stratified therapy in glioblastoma? *Dis. Markers* 2007;23:97-104.
- Hayashi K, Yoshida K, Matsui Y. A histone H3 methyltransferase controls epigenetic events required for meiotic prophase. *Nature* 2005; 438:374-8.
- Hebbes TR, Throne AW, Crane-Robinson C. A direct link between core histone acetylation and transcriptionally active chromatin. *EMBO J.* 1988;7:1395-402.
- Hede K. Histone deacetylase inhibitors sit at crossroads of diet, aging, cancer. *J. Natl. Cancer Inst.* 2006;98:377-9.
- Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, Kros JM, Hainfellner JA, Mason W, Mariani L, Bromberg JE, Hau P, Mirimanoff RO, Cairncross JG, Janzer RC, Stupp R. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N. Engl. J. Med.* 2005;352:997-1003.
- Hendzel MJ, Wei Y, Manchini MA. Mitosis-specific phosphorylation of histone H3 initiates primarily within pericentromeric heterochromatin during G2 and spreads in an ordered fashion coincident with mitotic chromosome condensation. *Chromosoma* 1997;106:348-60.
- Henry KW, Berger SL. Trans-tail modifications: wedge or bridge? *Nat. Struct. Biol.* 2002;9:565-6.
- Henry KW, Wyce A, Lo WS, Duggan LJ, Emre NCT, Kao CF, Pillus L, Shilatifard A, Osley MA, Berger SL. Transcriptional activation via sequential histone H2B ubiquitination and deubiquitination, mediated by SAGA-associated Ubp8. *Genes Dev.* 2003;17:2648-63.
- Herceg Z. Epigenetics and cancer: towards an evaluation of the impact of environmental and dietary factors. *Mutagenesis* 2007; 22:91-103.
- Herman JG, Graff JR, Myöhänen S, Nelkin BD, Baylin SB. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc. Natl. Acad. Sci. USA* 1996;93:9821-6.
- Hermann A, Gowher H, Jeltsch A. Biochemistry and biology of mammalian DNA methyltransferases. *Cell. Mol. Life Sci.* 2004;61:2571-87.
- Hoek HW, Brown AS, Susser E. The Dutch famine and schizophrenia spectrum disorders. *Soc. Psychiatry Psychiatr. Epidemiol.* 1998;33:373-9.

- Hoke SMT, Liang G, Mutis AI, Genereaux J, Brandi CJ. C-terminal processing of yeast Spt7 occurs in the absence of functional SAGA complex. *BMC Biochem.* 2007; 8:16.
- Holliday R, Pugh JE. DNA modification mechanisms and gene activity during development. *Science* 1975; 187:226-32.
- Holmstrom S, van Antwerp ME, Iñiguez-Lluhi JA. Direct and distinguishable inhibitory roles for SUMO isoforms in the control of transcriptional synergy. *Proc. Natl. Acad. Sci. USA* 2003;100:15758-63.
- Hong L, Schroth GP, Matthews HR, Yau P, Bradbury EM. Studies of the DNA binding properties of histone H4 therminus. Thermal denaturation studies reveal that acetylation markedly reduces the binding constant of the H4 "tail" to DNA. *J. Biol. Chem.* 1993;268:305-14.
- Horrobin DF. *The Madness of Adam and Eve. How Schizophrenia Shaped Humanity.* Berkshire (UK): Cox & Wyman Ltd Reading; 2001.
- Hsieh CL. In vivo activity of murine de novo methyltransferases, Dnmt3a and Dnmt3b. *Mol. Cell Biol.* 1999;19:8211-8.
- Hsiung DT, Marsit CJ, Houseman EA, Eddy K, Furniss CS, McClean MD, Kelsey KT. Global DNA methylation level in whole blood as a biomarker in head and neck squamous cell carcinoma. *Cancer Epidemiol. Biomarkers Prev.* 2007;16:108-14.
- Hsu DW, Lin MJ, Lee TL, Wen SC, Chen X, Shen CKJ. Two major forms of DNA (cytosine-5) methyltransferases in human somatic tissues. *Proc. Natl. Acad. Sci. USA* 1999;96:9751-6.
- Hsu JY, Sun, ZW, Li X, Reuben M, Tatchell K, Bishop DK, Grushcow JM, Brame CJ, Caldwell JA, Hunt DF, Lin R, Smith MM, Allis CD. Mitotic phosphorylation of histone H3 is governed by Ip1/aurora kinase and Glc7/PP1 phosphatase in budding yeast and nematodes. *Cell* 2000;102:279-91.
- Hu M, Yao J, Cai L, Bachman KE, van den Brûle F, Velculescu V, Polyak K. Distinct epigenetic changes in the stromal cells of breast cancers. *Nat. Genet.* 2005;37:899-905.
- Huang H, Kahana A, Gottschling DE, Prakash L, Lieberman SW. The ubiquitin-conjugating enzyme Rad6 (Ubc2) is required for silencing in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 1997;17:6693-9.
- Huang N, von Baur E, Garnier JM, Lerouge T, Vonesch JL, Lutz Y, Chambon P, Losson R. Two distinct nuclear receptor interaction domains in NSD1, a novel SET protein that exhibits characteristics of both corepressors and coactivators. *EMBO J.* 1998; 17:3398-412.
- Huang S. Histone methyltransferases, diet nutrients and tumour suppressors. *Nat. Rev. Cancer.* 2002;2:469-72.
- Huang SY, Barnard MB, Xu M, Matsui S, Rose SM, Garrard WT. The active immunoglobulin k chain gene is packaged by non-ubiquitin-conjugated nucleosomes. *Proc. Natl. Acad. Sci.* 1986;83:3738-42.
- Huang Y, Fang J, Bedford MT, Zhang Y, Xu RM. Recognition of histone H3 lysine-4 methylation by the double Tudor domain of JMJD2A. *Science* 2006;312:748-51.
- Hubbert C, Guardiola A, Shao R, Kawaguchi Y, Ito A, Nixon A, Yoshida M, Wang XF, Yao TP. HDAC6 is a microtubule-associated deacetylase. *Nature* 2002;417:455-8.
- Hughes C, Russell J, Robbins. TW. Evidence of executive dysfunction in autism. *Neuropsychologia* 1994; 32:477-92.
- Huyen Y, Zgheib O, Ditullio RA Jr, Gorgoulis VG, Zacharatos P, Petty TJ, Sheston EA, Mellert HS, Stavridi ES, Halazonetis TD. Methylated lysine 79 of histone H3 targets 53BP1 to DNA double-strand breaks. *Nature* 2004;432:406-11.
- Hwang WW, Venkatasubrahmanyam S, Ianculescu AG, Tong A, Boone C, Madhani HD. A conserved RING finger protein required for histone H2B monoubiquitination and cell size control. *Mol. Cell* 2003;11:261-6.
- Hymes J, Wolf B. Human biotinidase isn't just for recycling biotin. *J. Nutr.* 1999;129:485S-489S.
- Hymes J, Fleischhauer K, Wolf B. Biotinylation of histones by human serum biotinidase: assessment of biotinyl-transferase activity in sera from normal individuals and children with biotinidase deficiency. *Biochem. Mol. Med.* 1995;56:76-83.
- Iizuka M, Stillman B. Histone acetyltransferase HBO1 interacts with the ORC1 subunit of the human initiator protein. *J. Biol. Chem.* 1999;274:23027-34.

ISCID Encyclopedia of Science and Philosophy: hnRNA. International Society for Complexity, Information and Design. Disponible en: <http://www.iscid.org/encyclopedia/hnRNA>. Accesado el 16 de mayo 2008.

Issa JP. Epigenetic variation and human disease. *J. Nutr.* 2002;132:2388S-2392S.

Issa JP, Garcia-Manero G, Giles FJ, Mannari R, Thomas D, Faderl S, Bayar E, Lyons J, Rosenfeld CS, Cortes J and Kantarjian HM. Phase 1 study of low-dose prolonged exposures schedules of the hypomethylating agent 5-aza-2'-deoxycytidine (decitabine) in hematopoietic malignancies. *Blood* 2004;103:1635-40.

Ivanov D, Schleifer A, Eisenhaber F, Mechtler K, Haering T, Nasmyth K. Eco1 is a novel acetyltransferase that can acetylate proteins involved in cohesion. *Curr. Biol.* 2002;12:323-8.

Jackson JP, Lindroth AM, Cao X, Jacobsen SE. Control of CpNpG DNA methylation by the KRYPTONITE histone H3 methyltransferase. *Nature* 2002;416:556-60.

Jacob RA, Gretz DM, Taylor PC, James SJ, Pogribny IP, Miller BJ, Henningdagger SM, Swendseid ME. Moderate folate depletion increases plasma homocysteine and decreases lymphocyte DNA methylation in postmenopausal women. *J. Nutr.* 1998; 128:1204-12.

Jacobs JJ, van Lohuizen M. Polycomb repression: from cellular memory to cellular proliferation and cancer. *Biochim. Biophys. Acta* 2002; 1602:151-61.

Jacobs SA, Khorasanizadeh S. Structure of a HP1 chromodomain bound to a lisien 9-methylated histone H3 tail. *Science* 2002; 295:2080-3.

Jacobs SA, Harp JM, Devarakonda S, Kim Y, Rastinejad F, Khorasanizadeh S. The active site of the SET domain is construct on a knot. *Nat. Struct. Biol.* 2002; 9:833-8.

Jacobson EL, Antol KM, Juarez-Salinas H, Jacobson MK. Poly(ADP-ribose) metabolism in ultraviolet irradiated human fibroblasts. *J. Biol. Chem.* 1983; 258:103-7.

Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat. Genet.* 2003; 33(S):245-54.

Jain KK. Applications of biochips: from diagnostics to personalizad medicine. *Curr. Opin. Drug. Discov. Devel.* 2004;7:285-9.

James SJ, Cutler P, Melnyk S, Jernigan S, Janak L, Gaylor DW, Neubrander JA. 2004. Metabolic biomarkers of increased oxidative stress and impaired methylation capacity in children with autism. *Am. J. Clin. Nutr.* 2004;80:1611-7.

Jason LJ, Moore SC, Ausio J, Lindsey G. Magnesium-dependent association and folding of oligonucleosomes reconstituted with ubiquitininated H2A. *J. Biol. Chem.* 2001;276:14597-601.

Jeltsch A. Beyond Watson and Crick: DNA methylation and molecular enzymology of DNA methyltransferases. *Chembiochem* 2002;3:274-93.

Jentsch S, McGrath JP, Varshavsky A. The yeast DNA repair gene RAD6 encodes a ubiquitina-conjugating enzyme. *Nature* 1987;329:131-4.

Jenuwein T, Allis CD. Translating the histone code. *Science* 2001;293:1074-80.

Jeong JW, Bae MK, Ahn MY, Kim SH, Sohn TK, Bae MH, Yoo MA, Song EJ, Lee KJ, Kim KW. Regulation and destabilization of HIF-1alpha by ARD1-mediated acetylation. *Cell* 2003;111:709-20.

Jhaveri MS, Wagner C, Trepel JB. Impact of extracellular folate levels on global gene expression. *Mol. Pharmacol.* 2001;60:1288-95.

John RM, Surani MA. Agouti germ line gets acquisitive. *Nat. Genet.* 1999; 23:254-55.

Johnson ES, Gupta AA. An E3-like factor that promotes SUMO conjugation to the yeast septins. *Cell* 2001;106:735-44.

Johnstone RW. Histone deacetylase inhibitors: novel drugs for the treatment of cancer. *Nat. Rev. Drug Discovery* 2002;1:287-99.

Jones PA, Baylin SB. The fundamental role of epigenetic events in cancer. *Nat. Rev. Genet.* 2002;3:415-28.

Jones PA, Taylor SM. Cellular differentiation, cytidine analogs and DNA methylation. *Cell* 1980;20:85-93.

- Jose B, Oniki Y, Kato T, Nishino N, Sumida Y, Yoshida M. Novel histone deacetylase inhibitors: cyclic tetrapeptide with trifluoromethyl and pentafluoroethyl ketones. *Bioorg. Med. Chem. Lett.* 2004;14:5343-6.
- Juarez-Salinas H, Levi H, Jacobson EL, Jacobson MK. Poly(ADP-ribose) has a branched structure in vivo. *J. Biol. Chem.* 1982;257:607-9.
- Juarez-Salinas H, Mendoza-Alvarez H, Levi V, Jacobson MK, Jacobson EL. Simultaneous determination of linear and branched residues in poly(ADP-ribose). *Annals Biochem.* 1983;131:410-8.
- Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality determined by nutrition during parents' and grandparents' slow growth period. *Eur. J. Hum. Genet.* 2002; 10:682-8.
- Kagey MH, Mellhuish TA, Wotton D. The poly-comb protein Pc2 is a SUMO E3. *Cell* 2003;113:127-37.
- Kahyo T, Nishida T, Yasuda H. Involvement of PIAS1 in the sumoylation of tumor suppressors p53. *Mol. Cell* 2001;8:713-8.
- Kalebic T. Epigenetic changes: potential therapeutic targets. *Ann. N.Y. Acad. Sci.* 2003;983:278-85.
- Kamime J, Elangovan B, Subramanian T, Coleman D, Chinnadurai G. Identification of a cellular protein that specifically interacts with the essential cysteine region of the HIV-1 Tat transactivator. *Virology* 1996;216:357-66.
- Kaminker PG, Kim SH, Taylor RD, Zebarjadian Y, Funk WD, Morin GB, Yaswen P, Campisi J. TANK2, a new TRF1-associated polu(ADP-ribose) polymerase, causes rapad induction of cell death upon overexpression. *J. Biol. Chem.* 2001;276:35891-9.
- Kaminskas E, Farrell A, Abraham S, Baird A, Hsieh LS, Lee SL, Leighton JK, Patel H, Rahman A, Sridhara R, Wang YC, Pazdur R; FDA. Approval summary: azacitidine for treatment of myelodysplastic syndrome subtypes. *Clin. Cancer Res.* 2005;11:3604-8.
- Kaminsky Z, Wang SC, Petronis A. Complex disease, gender and epigenetics. *Ann. Med.* 2006; 38:530-44.
- Kanagawa T. Bias and artifacts in multitemplate polymerase chain reactions (PCR). *J. Biosci. Bioeng.* 2003;96:317-23.
- Kanduri C, Pant V, Loukinov D, Pugacheva E, Qi CF, Wolffe A, Ohlsson R, Lobanenkov VV. Functional association of CTCF with the insulator upstream of the H19 gene is parent of origin-specific and methylation sensitive. *Curr. Biol.* 2000;10:853-6.
- Kao CF, Hillyer C, Tsukuda T, Henry K, Berger S, Osley MA. Rad6 plays a role in transcriptional activation through ubiquitylation of histone H2B. *Genes Dev.* 2004;18:184-95.
- Kao HY, Lee CH, Komarov A, Han CC, Evans RM. Isolation and characterization of mammalian HDAC10, a novel histone deacetylase. *J. Biol. Chem.* 2002;277:187-93.
- Kaput J. Diet-disease gene interactions. *Nutrition* 2004; 20:26-31.
- Karamouzis MV, Konstantinopoulos PA, Papavassiliou AG. Roles of CREB-binding protein (CBP)/p300 in respiratory epithelium tumorigenesis. *Cell Res.* 2007;17:324-32.
- Karras GI, Kustatscher G, Buhecha HR, Allen MD, Pugieux C, Sait F, Bycroft M, Ladurner AG. The macro domain is an ADP-ribose binding module. *EMBO J.* 2005;24:1911-20.
- Kato Y, Salumbides BC, Wang XF, Qian DZ, Williams S, Wei Y, Sanni TB, Atadja P, Pili R. Antitumor effect to the histone deacetylase inhibitor LAQ824 in combination with 13-cis-retinoic acid in human malignant melanoma. *Mol Cancer Ther* 2007;6:70-81.
- Kawakami K, Brabender J, Lord RV, Groshen S, Greenwald BD, Krasna MJ, Yin J, Fleisher AS, Abraham JM, Beer DG, Sidransky D, Huss HT, Demeester TR, Eads C, Laird PW, Ilson DH, Kelsen DP, Harpole D, Moore MB, Danenberg KD, Danenberg PV, Meltzer SJ. Hypermethylated APC DNA in plasma and prognosis of patients with esophageal adenocarcinoma. *J. Natl. Cancer Inst.* 2000;92:1805-11.
- Kawasaki H, Schiltz L, Chiu R, Itakura K, Taira K, Nakatani Y, Yokoyama KK. ATF-2 has intrinsic histone acetyltransferase activity which is modulated by phosphorylation. *Nature* 2000;405:195-200.
- Kennedy BK, Barbie DA, Classon M, Dyson N, Harlow E. Nuclear organization of DNA replication in primary mammalian cells. *Genes Dev.* 2000;14:2855-68.

- Kersting M, Friedl C, Kraus A, Behn M, Pankow W, Schuermann M. Differential frequencies of p16(INK4a) promoter hypermethylation, p53 mutation, and K-ras mutation in exfoliative material mark the development of lung cancer in symptomatic chronic smokers. *J. Clin. Oncol.* 2000;18:3221-9.
- Kickhoefer VA, Siva AC, Kedersha NL, Inman EM, Ruland C, Streuli M, Rome LH. The 193-kD vault protein, VPARP, is a novel poly(ADP-ribose) polymerase. *J. Cell Biol.* 1999;146:917-28.
- Kijima M, Yoshida M, Sugita K, Horinouchi S, Beppu T. Trapoxin, an antitumor cyclic tetrapeptide, is an irreversible inhibitor of mammalian histone deacetylase. *J. Biol. Chem.* 1993;268:22429-35.
- Kim J, Cantwell CA, Johnson PF, Pfarr CM, Williams SC. Transcriptional activity of CCAAT/enhancer-binding proteins is controlled by a conserved inhibitory domain that is a target for sumoylation. *J. Biol. Chem.* 2002a; 277:38037-44.
- Kim J, Daniel J, Espejo A, Lake A, Krishna M, Xia L, Zhang Y, Bedford MT. Tudor, MBT and cromo domains gauge the degree of lisien methylation. *EMBO Rep.* 2006; 7:397-403.
- Kim KC, Huang S. Histone methyltransferases in tumor suppression. *Cancer Biol. Ther.* 2003; 2:491-9.
- Kim KC, Geng L, Huang S. Inactivation of a histone methyltransferase by mutations in human cancers. *Cancer Res* 2003a; 63:7619-23.
- Kim KI, Baek SH, Jeon YJ, Nishimori S, Suzuki T, Uchida S, Shimbara N, Saitoh H, Tanaka K, Chung CH.A new SUMO-1 specific protease, SUSP1, that is highly expressed in reproductive organs. *J. Biol. Chem.* 2000;275:14102-6.
- Kim M, Ahn JW, Song K, Paek KH, Pai HS. Forkhead-associated domains of the tobacco NtFHA1 transcription activator and the yeast Fh1 forkead transcription factor are functionally conserved. *J. Biol. Chem.* 2002b; 277:38781-90.
- Kim MY, Mauro S, Gevry N, Lis JT, Kraus WL. NAD<sup>+</sup>-dependent modulation of chromatin structure and transcription by nucleosome binding properties of PARP-1. *Cell* 2004a; 119:803-14.
- Kim SH, Jeong JWJ, Park JA, Lee JW, Seo JH, Jung BK, Bae MK, Kim KW. Regulation of the HIF-1a stability by histone deacetylases. *Oncol. Rep.* 2007; 17:647-51.
- Kim TG, Kraus JC, Chen J, Lee Y. JUMONJI, a critical factor for cardiac development, functions as a transcriptional repressor. *J. Biol. Chem.* 2003b; 278:42247-55.
- Kim TG, Chen J, Sadoshima J, Lee Y. Jumonji represses atrial natriuretic factor gene expression by inhibiting transcriptional activities of cardiac transcripcion factors. *Mol Cell Biol* 2004b; 24:10151-60.
- Kim YB, Lee KH, Sugita K, Yoshida M, Horinouchi S. Oxamflatin is a novel antitumor compound that inhibits mammalian histone deacetylase. *Oncogene* 1999; 18:2461-70.
- Kim YI. Folate and DNA methylation: a mechanistic link between folate deficiency and colorectal cancer? *Cancer Epidemiol. Biomarkers Prev.* 2004; 13:511-9.
- Klenova EM, Morse HC, Ohlsson R, Lovanekov VV. The novel BORIS + CTCF gene family is uniquely involved in the epigenetics of normal biology and cancer. *Sem. Canc. Biol.* 2002;449:1-16.
- Klose RJ, Yamane K, Bae Y, Zhang D, Erdjument-Bromage H, Tempst P, Wong J, Zhang Y. The transcriptional repressor JHDM3A demethylates trimethyl histone H3 lysine 9 and lisien 36. *Nature* 2006;442:312-6.
- Klymenko T, Papp B, Fischle W, Köcher T, Schelder M, Fritsch C, Wild B, Wilm M, Müller J. A polycomb group protein complex with sequence- specific DNA-binding and selective methyl-lysine-binding activities. *Genes Dev.* 2006;20:1110-22.
- Kobza K, Camporeale G, Rueckert B, B, Kueh A, Griffin JB, Sarath G, Zempleni J. K4, K9 and K18 of human histone H3 are targets for biotinylation by biotinidase. *FEBS J.* 2005;272:4249-59.
- Koch-Nolte F, Haag F. Mono(ADP-ribosyl)transferases and related enzymes in animal tisúes. Emerging gene families. *Adv. Exp. Med. Biol.* 1997;419:1-13
- Kohlmaier A, Savarese F, Lachner M, Martens J, Jenuwein T, Wultz A. A crhomosomal memory triggered by Xist regulates histone methylation in X inactivation. *PLoS Biol.* 2004;2:991-1003.

- Koken MH, Reynolds P, Jaspers-Dekker I, Prakash L, Prakash S, Bootsma D, Hoeijmakers JH. Structural and functional conservation of two human homologs of the yeast DNA repair gene RAD6. *Proc. Natl. Acad. Sci.* 1991;88:8865-9.
- Komite C, Watanabe T, Katayama Y, Yoshino A, Yokohama T, Fukushima T. Promoter hypermethylation of the DNA repair gene O6-methylguanine-DNA methyltransferase is an independent predictor of shortened progression free survival in patients with low-grade diffuse astrocytomas. *Brain Pathol.* 2003;13:176-84.
- Konings EJ, Goldbohm RA, Brants HA, Saris WH, van den Brandt PA. Intake of dietary folate vitamers and risk of colorectal carcinoma; results from The Netherlands Cohort Study. *Cancer* 2002;95:1421-33.
- Kornman K, Fogarty C, Mezkin MS, Bidlack WR, Randolph RK(eds). *Nutrigenomics: Opportunities and Challenges. Phytochemicals: Nutrient-Gene Interaction*. Boca Ratón (EUA): Taylor & Francis; 2006.
- Kotaja N, Aittomaki S, Silvennoinen O, Palvinmo JJ, Janne OA. ARIP3 (androgen receptor-interacting protein 3) and other PIAS (protein inhibitor of activated STAT) proteins differ in their ability to modulate steroid receptor-dependent transcripcional activation. *Mol. Endocrinol.* 2000;14:1986-2000.
- Kothapalli N, Zempleni J. Double strand breaks of DNA decrease biotinylation of lysine-12 in histone H4 in JAR cells. *FASEB J.* 2004;18:A103-4.
- Kothapalli N, Zempleni J. Biotinylation of histones depends on the cell cycle in NCI-H69 small cell lung cancer cells. *FASEB J* 2005;19:A55.
- Kothapalli N, Camporeale G, Kueh A, Chew YC, Oommen AM, Griffin JA, Zempleni J. Biological functions of biotinylated histones. *J Nutr Biochem* 2005a;16:446-8.
- Kothapalli N, Sarath G, Zempleni J. Biotinylation of K12 in histone H4 decreases in response to DNA double stand breaks in human Jar choriocarcinoma cells. *J. Nutr.* 2005b;135:2337-42.
- Kouzarides T. Acetylation: a regulatory modification to rival phosphorylation? *EMBO J.* 2000; 19:1176-9.
- Kouzarides T. Histone methylation in transcriptional control. *Curr. Opin. Genet. Dev.* 2002;12:198-209.
- Kouzarides T. Chromatin modifications and their function. *Cell* 2007; 128:693-705.
- Kreimeyer A, Wielckens K, Adamietz P, Hilz H. DNA repair-associated ADP-ribosylation in vivo. Modification of histone H1 differs from that of the principal acceptor proteins. *J. Biol. Chem.* 1984;259:890-6.
- Kreimeyer A, Adamietz P, Hilz H. Alkylation-induced mono-(ADP-ribosyl)-histones H1 and H2B. Hydroxylamine-resistant linkage in hepatoma cells. *Biol. Chem. Hoppe Seyler* 1985;366:537-44.
- Krogan NJ, Kim M, Tong A, Golshani A, Cagney G, Canadian V, Richards DP, Beattie BK, Emili A, Boone C, Shilatifard A, Buratowski S, Greenblatt J. Methylation of histone H3 by Set2 in *Saccharomyces cerevisiae* is linked to transcriptional elongation by RNA polymerase II. *Mol. Cell Biol.* 2003a;23:4207-18.
- Krogan NJ, Dover J, Wood A, Schneider J, Heidt J, Boateng MA, Dean K, Ryan OW, Golshani A, Johnston M, Greenblatt JF, Shilatifard A. The Paf1 complex is required for histone H3 methylation by COMPASS and Dot1p: linking transcriptional elongation to histone methylation. *Mol. Cell* 2003b;11:721-9.
- Kuendgen A, Knipp S, Fox F, Strupp C, Hildebrandt B, Steidl C, Germing U, Haas R, Gattermann N. Results of a phase 2 study of valproic acid alone or in combination with all-trans retinoic acid in 75 patients with myelodysplastic syndrome and relapsed or refractory acute myeloid leukemia. *Ann. Hematol.* 2005;84:61-6.
- Kustatscher G, Hothorn M, Pugieux C, Schelfzek K, Ladurner AG. Splicing regulates NAD metabolite binding to histone macro H2A. *Nat. Struct. Mol. Biol.* 2005;12:624-5.
- Kuzmichev A, Jenuwein T, Tempst P, Reinberg D. Different EZH2-containing complexes target methylation of histone H1 or nucleosomal histone H3. *Mol. Cell* 2004;14:183-93.
- Kwak YT, Guo J, Prajapati S, Park KJ, Surabhi RM, Miller B, Gehrig P, Gaynor RB. Methylation of SPT5 regulates its interaction with RNA polymerase II and transcriptional elongation properties. *Mol. Cell* 2003;11:1055-66.
- Kwon T, Chang JH, Kwak E, Lee CW, Joachimiak A, Kim YC, Lee J, Cho Y. Mechanism of histone lysine methyl transfer revealed by the structure of SET7/9-AdoMet. *EMBO J.* 2003;22:292-303.
- Lachner M, O'Carroll D, Rea S, Mechtler K, Jenuwein T. Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature* 2001;410:116-20.

- Lachner M, O'Sullivan RJ, Jenuwein T. An epigenetic road map for histone lisien methylation. *J. Cell. Sci.* 2003;116:2117-24.
- Lacoste N, Ultey RT, Hunter JM, Poirier GG, Cote J. Disruptor of telomeric silencing-1 is a chromatin-specific histone H3 methyltransferase. *J. Biol. Chem.* 2002;277:30421-4.
- Lagger G, O'Carrol D, Rembold M, Khier H, Tischler J, Weitzer G, Schuettengruber B, Hauser C, Brunmeir R, Jenuwein T, Seiser C. Essential function of histone deacetylase 1 in proliferation control and CDK inhibitor repression. *EMBOJ.* 2002;21:2672-81.
- Laird PW. The power and the promise of DNA methylatio markers. *Nat. Rev. Cancer* 2003;3:253-66.
- Laird PW. Cancer epigenetics. *Hum. Mol. Genet.* 2005;14:R65-T76.
- Laird PW, Jackson-Grusby L, Fazeli A, Dickinson SL, Jung WE, Li E, Weinberg RA, Jaenisch R. Supression of intestinal neoplasia by DNA hypomethylation. *Cell* 1995;81:197-205.
- Lamb JA, Banby G, Bonora E, Sykes N, Bacchelli E, Blasi F, Maestrini E, Broxholme J, Tzenova1 J, Weeks D, Bailey AJ, Monaco AP. Análisis of IMGSAC autism susceptibility loci: evidence for sex limited and parent of origin specifi effects. *J. Med. Genet.* 2005;42:132-37.
- Lamming DW, Latorre-Esteves M, Medvedik O, Wong SN, Tsang FA, Wang C, Lin SJ, Sinclair DA. HST2 mediates SIR2-independent life-span extensión by calorie restriction. *Science* 2005;309:1861-4.
- Lanctôt C, Cheutin T, Cremer M, Cavalli G, Cremer T. Dynamic genome architecture in the nuclear space : regulation of gene expression in three dimensions. *Nat. Rev. Genet.* 2007;8:104-15.
- Lauster R, Trautner TA, Noyer-Weidner M. Cytosine-specific type II DNA methyltransferases: a conserved enzyme core with variable target-recognizing domains. *J. Mol. Biol.* 1989;206:305-12.
- Le Bivic A, Hirn M, Reggio H. HT-29 cells are an in vitro model for the generation of cell polarity in epithelia during embryonic differentiation. *PNAS* 1988; 85:136-40.
- Le Drean Y, Mincheneau N, Le Goff P, Michel D. Potentiation of glucocorticoid receptor transcriptional activity by sumoylation. *Endocrinology* 2002;143:3482-9.
- Leduc Y, de Murcia G, Lamarre D, Poirier GG. Visualization of poly(ADP-ribose) synthetase associated with polynucleosomes by immunoelectron microscopy. *Biochim. Biophys. Acta* 1986;885:248-55.
- Lee BH, Yegnasubramanian S, Lin X, Nelson WG. Procainamide is a specific inhibitor of DNA methyltransferase 1. *J. Biol. Chem.* 2005a;280:40749-56.
- Lee DY, Teyssier C, Strahl BD, Stallcup MR. Role of protein methylation in regulation of transcription. *Endocr. Rev.* 2004;2:147-70.
- Lee JH, Park JH, Jung Y, Kim JH, Jong HS, Kim TY, Bang YJ. Histone deacetylase inhibitor enhances 5-fluorouracil cytotoxicity by down-regulating thymidylate synthase in human cancer cells. *Mol. Cancer Ther.* 2006;5:3085-95.
- Lee JW, Choi HS, Gyuris J, Brent R, Moore DD. Two classes of proteins dependent on either the presence or absence of thyroid hormone for interaction with the thyroid hormone receptor. *Mol. Endocrinol.* 1995; 9:243-54.
- Lee SH, Oshige M, Durant ST, Rasila KK, Williamson EA, Ramsey H, Kwan L, Nickoloff JA, Hromas R. The SET domain protein Metnase mediates foreign DNA integration and links integration to nonhomologous end-joining repair. *Proc. Natl. Acad. Sci. USA* 2005b; 102:18075-80.
- Lee MG, Wynder C, Cooch N, Shiekhattar R. An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* 2005c;437:432-5.
- Lehnertz B, Ueda Y, Derikck AA, Braunschweig U, Perez-Burgos L, Kubicek S, Chen T, Li E, Jenuwein T, Peters AH. Suv39h-mediated histone H3 lysine 9 methylation directs DANN methylation to major satellite repeats at pericentric heterochromatin. *Curr. Biol.* 2003;12:1192-200.
- Lei H, Oh SP, Okano M, Jüttermann R, Goss KA, Jaenisch R, Li E. De novo DNA cytosine methyltransferase activities in mouse embryonic stem cells. *Development* 1996;122:3195-205.

- Lemieux ME, Yang X, Jardine K, He X, Jacobsen KX, Staines WA, Harper ME, McBurney MW. The SIRT1 deacetylase modulates the insulin-like growth factor signaling pathway in mammals. *Mech. Ageing Dev.* 2005;126:1097-105.
- Leonhardt H, Page AW, Weier HU, Bestor TH. A targeting sequence directs DNA methyltransferase to sites of DNA replication in mammalian nuclei. *Cell* 1992;71:865-73.
- Leuwen F van, Grafken PR, Gottschling DE. Dot1p modulates silencing in yeast by methylation of nucleosome core. *Cell* 2002;109:745-56.
- Levinger L, Vashavsky A. Selective arrangement of ubiquitinated and D1 protein-containing nucleosomes within the *Drosophila* genome. *Cell* 1982;278:8897-903.
- Lewis A, Reik W. How imprinting centres work. *Cytogenet. Genome Res.* 2006;113:81-9..
- Li E. Chromatin modification and epigenetic reprogramming in mammalian development. *Nat. Rev. Genet.* 2002a; 3:662-73.
- Li B, Howe L, Anderson S, Yates JR III, Workman JL. The Set2 histone methyltransferase functions through the phosphorylated carboxy-terminal domain of RNA polymerase II. *J. Biol. Chem.* 2003;278:8897-903.
- Li E, Bestor TH, Jaenisch R. Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* 1992;69:915-26.
- Li E, Beard C, Jaenisch R. Role for DNA methylation in genomic imprinting. *Nature* 1993; 366:362-5.
- Li H, Leo C, Zhu J, Wu X, O'Neil J, Park EJ, Chen JD. Sequestration and inhibition of Daxx-mediated transcriptional repression by PML. *Mol. Cell Biol.* 2000; 20:1784-96.
- Li H, Rauch T, Chen ZX, Szabó PE, Riggs AD, Pfeiffer GP. The histone methyltransferase SETDB1 and the DNA methyltransferase DNMT3A interact directly and localize to promoters silenced in cancer cells. *J. Biol. Chem.* 2006a; 281:19489-500.
- Li H, Ilin S, Wang W, Duncan EM, Wysocka J, Allis CD, Patel DJ. Molecular basis for site-specific read-out of histone H3K4me3 by the BPTF PHD finger of NURF. *Nature* 2006b; 442:91-5.
- Li J, Moazed D, Gygi SP. Association of the histone methyltransferase Set2 with RNA polymerase II plays a role in transcription elongation. *J. Biol. Chem.* 2002b; 277:49383-88.
- Li SJ, Hochstrasser M. A new protease required for cell-cycle progression in yeast. *Nature* 1999; 398:246-51.
- Liang H, Ward WF. PCG-1a: a key regulation of energy metabolism. *Advan. Physiol. Edu.* 2006;30:145-51.
- Lifton RP, Goldberg ML, Karp RW, Hogness DS. The organization of the histone genes in *Drosophila melanogaster*: functional and evolutionary implications. *Cold Spring Harb. Symp. Quant. Biol.* 1978; 42:1047-51.
- Liszt G, Ford E, Kurtev M, Guarente L. Mouse Sir2 homolog SIRT6 is a nuclear ADP-ribosyltransferase. *J. Biol. Chem.* 2005;280:21313-20.
- Liu Y, Oakeley EJ, Sun L, Jost J-P. Multiple domains are involved in the targeting of the mouse DNA methyltransferase to the DNA replication foci. *Nucleic Acids Res.* 1998;26:1038-45.
- Lizst G, Ford E, Kurtev M, Guarente L. Mouse Sir2 homolog SIRT6 is a nuclear ADP-ribosyltransferase. *J. Biol. Chem.* 2005;280:21313-20.
- Lo WS, Triev RC, Rojas JR, Duggan L, Hsu JY, Allis CD, Marmorstein R, Berger SL. Phosphorylation of serine 10 in histone H3 is functionally linked *in vitro* and *in vivo* to Gcn5-mediated acetylation at lysine 14. *Mol. Cell* 2000;5:917-26.
- Loprevite M, Tiseo M, Grossi F, Scolaro T, Semino C, Pandolfi A, Favoni R, Ardizzone A. In vitro study of CI-994, a histone deacetylase inhibitor, in non-small cell lung cancer cell lines. *Oncol. Res.* 2005;15:39-48.
- Lu Q, Wu A, Richardson BC. Demethylation of the same promoter sequence increases CD70 expression in lupus T cells and T cells treated with lupus-inducing drugs. *J. Immunol.* 2005; 174:6212-9.
- Lu Q, Qiu X, Hu N, Wen H, Su Y, Richardson BC. Epigenetics, disease and therapeutic interventions. *Aging Res. Rev.* 2006; 5:449-67.
- Lu ZH, Xu H, Leno GH. DNA replication in quiescent cell nuclei: regulation by the nuclear envelope and chromatin structure. *Mol. Biol. Cell* 1999; 10:4091-106.

- Lumey LH. Decreased birthweights in infants after maternal in utero exposure to the Dutch famine of 1944-1945. *Paediatr. Perinat. Epidemiol.* 1992; 6:240-53.
- Lund AH, van Lohuizen M. Epigenetics and cancer. *Genes Dev.* 2004;18:2315-35.
- Lund G, Anderson L, Lauria M, Lindholm M, Fraga MF, Villar-Garea A, Ballestar E, Esteller M, Zaina S. DNA methylation polymorphisms precede any histological sign of atherosclerosis in mice lacking apolipoprotein E. *J. Biol. Chem.* 2004;279:29147-54.
- Luo J, Nikolaev AY, Imai S, Chen D, Su F, Shiloh A, Guarente L, Gu W. Negative control of p53 by Sir2a promotes cell survival under stress. *Cell* 2001;107:137-48.
- Luo RX, Postigo AA, Deand DC. Rb interacts with histone deacetylase to repress transcription. *Cell* 1998; 92:463-73.
- Lyko F, Brown R. DNA methyltransferase inhibitors and the development of epigenetic cancer therapies. *J. Natl. Cancer Inst.* 2005;97:1498-506.
- Lyko F, Ramsahoye BH, Kashevsky H, Tudor M, Mastrangelo MA, Orr-Weaver TL, Jaenisch R. Mammalian (cytosine-5) methyltransferases cause genomic DNA methylation and lethality in Drosophila. *Nature Genet* 1999;23:363-6.
- Lyn-Cook BD, Blann E, Payne PW, Bo J, Sheehan D, Medlock K. Methylation profile and amplification of proto-oncogenes in rat pancreas induced with phytoestrogens. *Proc. Soc. Exp. Biol. Med.* 1995;208:116-9.
- Ma H, Baumann CT, Li H, Strahl BD, Rice R, Jelinek MA, Aswad DW, Allis CD, Hager GL, Stallcup MR. Hormone-dependent, CARM1-directed, arginina-specific methylation of histone H3 on a steroid-regulated promoter. *Curr Biol* 2001a;11:1981-5.
- Ma Q, Baldwin KT, Renzelli AJ, McDaniel A, Dong L. TCDD-inducible poly(ADP-ribose) polymerase: a novel response to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Biochem. Biophys. Res. Commun.* 2001b;289:499-506.
- MacGregor PF, Squire JA. Application of microarrays to the analysis of gene expression in cancer. *Clin. Chem.* 2002;48:351-8.
- MacLennan NK, James SJ, Melnyk S, Piroozi A, Jernigan S, Hsu JL, Janke SM, Pham TD, Lane RH. Uteroplacental insufficiency alters DNA methylation, one-carbon metabolism, and histone acetylation in IUGR rats. *Physiol Genomic* 2004;18:43-50.
- Magnaghi-Jaulin L, Groisman R, Naguibneva I, Robin P, Lorain S, Le Villain JP, Troalen F, Trouche D, Hareló-Bellan A. protein represses transcripción by recruiting a histone deacetylase. *Nature* 1998;391:601-5.
- Main ERG, Stott K, Jacksin SE, Regan L. Local and long-range stability in tandemly arrayed tetratricopeptide repeats. *Proc. Natl. Acad. Sci. USA* 2005;102:5721-6.
- Mak W, Nesterova TB, de Napoles M, Appanah R, Yamanaka S, Otte AP, Brockdorff N. Reactivation of the paternal X chrosomome in early mouse embryos. *Science* 2004;303:666-9.
- Manthey KC, Griffin JB, Zempleni J. Biotin supply affects expression of biotin transporters, biotinylation of carboxylases, and metabolism of interleukin-2 in Jurkat cells. *J. Nutr.* 2002;132:887-92.
- Mantripraga KK, Buckley PG, de Stahl TD, Dumanski JP. Genomic microarrays in the spotlight. *Trends Genet.* 2004;20:87-94.
- Marcucci G, Silverman L, Eller M, Lintz L, Beach CL. Bioavailability of azacitidine subcutaneous versus intravenous in patients with the myelodysplastic syndromes. *J. Clin. Pharmacol.* 2005;45:597-602.
- Margueron R, Trojer P, Reinberg D. The key to development: interpreting the histone code? *Curr. Opin. Genet. Dev.* 2005; 15:163-76.
- Mariadson JM. HDACs and HDAC inhibitors in colon cancer. *Epigenetics* 2008;3:28-37.
- Marino-Ramirez L, Spouge JL, Kanga GC, Landsman D. Statistical analysis of over-represented words in human promoter sequences. *Nucleic Acids Res.* 2004;32:949-58.
- Marks PA, Breslow R. Dimethyl sulfoxide to vorinostat: development of this histone deacetylase inhibitor as an anticancer drug. *Nat. Biotechnol.* 2007;25:84-90.
- Marmostein R. Structure of histone acetyltransferases. *J. Mol. Biol.* 2001;311:433-44.

- Martínez E, Palhan VB, Tjernberg A, Lymar ES, Gamper AM, KunduTK, Chait BT, Roeder RG. Human STAGA complex is a chromatin-acetylating transcription coactivator that interacts with pre-mRNA splicing and DNA damage-binding factors *in vivo*. *Mol. Cell Biol.* 2001;21:6782-95.
- Mathers JC, Hesketh JE. The biological revolution: understanding the impact of SNPs on diet-cancer interrelationships. *J. Nutr.* 2007; 137:253S-238S.
- McBride AE, Silver PA. State of the Arg: protein methylation at arginine comes of age. *Cell* 2001;106:5-8.
- McBurney MW, Yang X, Jardine K, Hixon M, Boekelheide K, Webb JR, Lansdorp PM, Lemieux M. The mammalian SIR2a protein has a role in embryogenesis and gametogenesis. *Mol. Cell Biol.* 2003;23:38-54.
- McKinsey TA, Zhang CL, Lu J, Olson EN. Signaling dependent nuclear export of a histone deacetylase regulates muscle differentiation. *Nature* 2000a; 408:106-11.
- McKinsey TA, Zhang CL, Olson EN. Activation of the myocyte enhancer factor-2 transcription factor by calcium/calmodulin-dependent protein kinase-stimulated binding 14-3-3 to histone deacetylase 5. *Proc. Natl. Acad. Sci. USA*. 2000b; 97:14400-5.
- McLeod KI, Goldrick RB, Whyte HM. The effect of maternal malnutrition on the progeny in the rat. Studies on growth, body composition and organ cellularity in first and second generation progeny. *Aust. J. Exp. Biol. Med.* 1972; 50:435-46.
- MedlinePlus. Síndrome metabólico. Disponible en: <http://www.nlm.nih.gov/medlineplus/spanish/metabolicsyndrome.html>. Accesado el 4 de junio 2008.
- Melchior F. SUMO-nonclassical ubiquitin. *Annu. Rev. Cell Dev. Biol.* 2000;16:591-626.
- Melzner I, Scott V, Dorsch K, Fischer P, Wabitsch M, Brüderlein S, Hasel C, Möller P. Leptin gene expression in human preadipocytes is switched on by maturation-induced demethylation of distinct CpGs in its proximal promoter. *J. Biol. Chem.* 2002;277(47):45420-7.
- Meng G, Inazawa J, Ishida R, Tokura K, Nakahara K, Aoki K, Kasai M. Structural analysis of the gene encoding RP58, a sequence-specific transrepressor associated with heterochromatin. *Gene* 2000;242:59-64.
- Menvielle G, Luce D, Goldberg P, Bugel I, Leclere A. Smoking, alcohol drinking and cancer risk for various sites of the larynx and hypopharynx. A case-control study in France. *Eur. J. Cancer Prev.* 2004;13:165-72.
- Merrow M, Spoelstra K, Roenneberg T. The circadian cycle: daily rhythms from behaviour to genes. *EMBO Rep.* 2005; 6:930-5.
- Metzger E, Wissmann M, Yin N, Müller JM, Schneider R, Peters AH, Günther T, Buettner R, Schüle R. LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature* 2005;437:436-9.
- Meyer-Ficca ML, Meyer RG, Coyle DL, Jacobson EL, Jacobson MK. Human poly(ADP-ribose) glycohydrolase is expressed in alternative splice variants yielding isoforms that localize to different cell compartments. *Exp. Cell Res.* 2004;297:521-32.
- Michan S, Sinclair D. Sirtuins in mammals: insights into their biological function. *Biochem. J.* 2007;404:1-13.
- Michishita E, Park JY, Burneskis JM, Barrett JC, Horikawa I. Evolutionary conserved and nonconserved cellular localizations and functions of human SIRT proteins. *Mol. Biol. Cell* 2005;16:4623-35.
- Miller JW, Nadeau MR, Smith J, Smith D, Selhub J. Folate-deficiency-induced homocysteinaemia in rats: disruption of S-adenosylmethionine's co-ordinate regulation of homocysteine metabolism. *Biochem. J.* 1994;298:415-9.
- Milne TA, Briggs SD, Brock HW, Martin ME, Gibbs D, Allis CD, Hess JL. MLL targets SET domain methyltransferase activity to Hox gene promoters. *Mol. Cell* 2002;10:1107-17.
- Milne TA, Hughes CM, Lloyd R, Yang Z, Rozenblatt-Rosen O, Dou Y, Schnepf RW, Krinkel C, Livolsi VA, Gibbs D, Hua X, Roeder RG, Meyerson M, Hess JL. Menin and MLL cooperatively regulate expression of cyclin-dependent kinase inhibitors. *Proc. Natl. Acad. Sci. USA* 2005;102:749-54.
- Min J, Zhang X, Cheng X, Grewal SI, Xu RM. Structure of the SET domain histone lysine methyltransferase Clr4. *Nat. Struct. Biol.* 2002;9:828-32.

- Min J, Zhang Y, Xu RM. Structural basis for specific binding of polycomb chromodomain to histone H3 methylated at Lys 27. *Genes Dev.* 2003a;17:1823-8.
- Min J, Feng Q, Li Z, Zhang Y, Xu RM. Structure of the catalytic domain of human DOT1L, a non-SET domain nucleosomal histone methyltransferase. *Cell* 2003b;112:711-23.
- Minaga T, Kun E. Spectral analysis of the conformation of polyadenosine diphosphoribose. Evidence indicating secondary structure. *J. Biol. Chem.* 1983;258:725-30.
- Mitsiades CS, Mitsiades NS, McMullan CJ, Poulaki V, Shringarpure R, Hideshima T, Akiyama M, Chauhan D, Munshi N, Gu X, Bailey C, Joseph M, Libermann TA, Richon VM, Marks PA, Anderson KC. Transcriptional signature of histone deacetylase inhibition in multiple myeloma: biological and clinical implications. *Proc. Natl. Acad. Sci. USA* 2004;101:540-5.
- Moazed D, Johnson D. A deubiquitinating enzyme interacts with SIR4 and regulates silencing in *S. cerevisiae*. *Cell* 1996;86:667-77.
- Moore SC, Jason L, Ausio J. The elusive structural role of ubiquitinated histones. *Biochem. Cell Biol.* 2002;80:311-9.
- Moraes RC, Blonder A, Birkenkamp-Demtroeder K, Tirard J, Orntoft TF, Gertler A, Durand P, Naville D, Bégeot M. Study of the alteration of gene expression in adipose tissue of diet-induced obese mice by microarray and reverse transcription-polymerase chain reaction analyses. *Endocrinology* 2003;144:4773-82.
- Morison IA, Ramsay JP, Spencer HG. A census of mammalian imprinting. *Trends Genet.* 2005;21:457-65.
- Mostoslavsky R, Chua KF, Lombard DB, Pang WW, Fischer MR, Gellon L, Liu P, Mostoslavsky G, Franco S, Murphy MM, Mills KD, Patel P, Hsu JT, Hong AL, Ford E, Cheng HL, Kennedy C, Nunez N, Bronson R, Frendewey D, Auerbach W, Valenzuela D, Karow M, Hottiger MO, Hursting S, Barrett JC, Guarente L, Mulligan R, Demple B, Yancopoulos GD, Alt FW. Genomic instability and aging-like phenotype in the absence of mammalian SIRT6. *Cell* 2006;124:315-29.
- Motamedi MR, Verdel A, Colmenares SU, Gerber SA, Gygi SP, Moazed D. Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell* 2004;119:789-802.
- Motta MC, Divecha N, Lemieux M, Kamel C, Chen D, Gu W, Bultsma Y, McBurney M, Guarente L. Mammalian SIRT1 represses forkhead transcription factors. *Cell* 2004;116:551-63.
- Moulton T, Crenshaw T, Hao Y, Moosikaswan J, Lin N, Dembitzer F, Hensle T, Weiss L, McMorrow L, Loew T, Kraus W, Gerald W, Tycko B. Epigenetic lesions at the H19 locus in Wilms' tumor patients. *Nature Genet.* 1994;7:440-7.
- Mujtaba S, He Y, Zeng S, Plotnikova O, Sachidanand Sanchez R, Zeleznik-Le NJ, Ronai Z, Zhou MM. Structural mechanism of the bromodomain of the coactivator CBP in p53 transcriptional activation. *Mol Cell* 2004;13:251-63.
- Müller J, Hart CM, Francis NJ, Vargas ML, Sengupta A, Wild B, Miller EL, O'Connor MB, Kingston RE, Simon JA. Histone methyltransferase activity of a Drosophila polycomb group repressor complex. *Cell* 2002;111:197-208.
- Müller M, Kersten S. Nutrigenomics: goals and strategies. *Nat. Rev. Genet.* 2003;4:315-22.
- Müller S, Hoege C, Pyrowolakis G, Jentsch S. SUMO, ubiquitin's mysterious cousin. *Nat. Rev. Mol. Cell. Biol.* 2001;2:202-10.
- Multinovic S, D'Alessio AC, Detich N, Szyf M. Valproate induces widespread epigenetic reprogramming which involves demethylation of specific genes. *Carcinogenesis* 2007;28:560-71.
- Munaka M, Kohshi K, Kawamoto T, Takasawa S, Nagata N, Itoh H, Oda S, Katoh T. Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and the risk of hepatocellular carcinoma. *J. Cancer Res. Clin. Oncol.* 2003;129:355-60.
- Murcia G de, Huletsky A, Lamarre D, Gaudreau A, Pouyet J, Daune M, Poirier GG. Modulation of chromatin superstructure induced by poly(ADP-ribose) synthesis and degradation. *J. Biol. Chem.* 1986;261:7011-17.
- Muskiet FA, Kemperman RF. Folate and long chain polyunsaturated fatty acids in psychiatric disease. *J. Nutr. Biochem.* 2006;17:717-27.

- Myers PZ. Basics: what is a gene? Pharyngula. 2007. Disponible en: [http://scienceblogs.com/pharyngula/2007/01/basics\\_what\\_is\\_a\\_gene.php](http://scienceblogs.com/pharyngula/2007/01/basics_what_is_a_gene.php). Accesado el 29 de mayo 2008.
- Myzak MC, Karphus PA, Chung FL, Dashwood RH. A novel mechanism of chemoprotection by sulphoraphane: Inhibition of histone deacetylase. *Cancer Res.* 2004;64:5767-74.
- Myzak MC, Ho E, Dashwood RH. Dietary agents as histone deacetylase inhibitors. *Mol. Carcinog.* 2006;45:443-6.
- Nakamura T, Blechman J, Tada S, RozovskaiaDagger T, Itoyama T, Bullrich F, Mazo A, Croce CM, GeigerDagger B, Canaani E. huASH1 protein, a putative transcription factor encoded by a human homologue of the Drosophila ash1 gene, localizes to both nuclei and cell-cell tight junctions. *Proc. Natl. Acad. Sci. USA* 2000;97:7284-9.
- Nakamura T, Mori T, Tada S, Krajewski W, Rozovskaia T, Wassell R, Dubois G, Mazo A, Croce CM, Canaani E. ALL-1 is a histone methyltransferase that assembles a supercomplex of proteins involved in transcriptional regulation. *Mol. Cell* 2002;10:1119-28.
- Nakatani Y. Histone acetylases-versatile players. *Genes Cell* 2001;6:79-86.
- Napoles M de, Mermoud JE, Wakao R, Tang YA, Endoh M, Appanah R, Nesterova TB, Silva J, Otte AP, Vidal M, Koseki H, Brockdorff N. Polycomb group proteins Ring1A/B link ubiquitylation of histone H2A to heritable gene silencing and X inactivation. *Dev. Cell* 2004;7:663-76.
- Narang MA, Dumas R, Ayer LM, Gravel RA. Reduced histone biotinylation in multiple carboxylase deficiency patients: a nuclear role for holocarboxylase synthetase. *Hum. Mol. Genet.* 2004;13:15-23.
- Nebbioso A, Clarke N, Voltz E, Germain E, Ambrosino C, Bontempo P, Alvarez R, Schiavone EM, Ferrara F, Bresciani F, Weisz A, de Lera AR, Gronemeyer H, Altucci L. Tumor-selective action of HDAC inhibitors involves TRAIL induction in acute myeloid leukemia cells. *Nat. Med.* 2005;11:77-84.
- Neel JV. Diabetes mellitus: a "thrifty" genotype rendered detrimental by "progress"? *Am. J. Hum. Genet.* 1962;14:353-62.
- Net Industries Science Encyclopedia. Disponible en: <http://science.jrank.org/pages/7435/Xenogamy.html>. Accesado el 1 de junio 2008.
- Neuwald AF, Landsman D. GCN5-related histone N-acetyltransferases belong to a diverse superfamily that includes the yeast SPT10 protein. *Trends Biochem.* 1997;22:154-5.
- Ng HH, Bird A. DNA methylation and chromatin modification. *Curr. Opin. Genet. Dev.* 1999; 9:158-63.
- Ng HH, Xu RM, Zhang Y, Struhl K. Ubiquitination of histone H2B by Rad6 is required for efficient Dot1-mediated methylation of histone H3 lysine 79. *J. Biol. Chem.* 2002a; 277:34655-7.
- Ng HH, Feng Q, Wang H, Erdjument-Bromage H, Tempst P, Zhang Y, Struhl K. Lysine methylation within the globular domain of histone H3 by Dot1 is important for telomeric silencing and Sir protein association. *Genes Dev.* 2002b; 16:1518-27.
- Ng HH, Robert F, Young RA, Struhl K. Targeted recruitment of Set1 histone methylase by elongating Pol II provides a localized mark and memory of recent transcriptional activity. *Mol Cell* 2003a; 11:709-19.
- Ng HH, Ciccone DN, Morshead KB, Oettinger MA, Struhl K. Lysine-79 of histone H3 is hypomethylated at silenced loci in yeast and mammalian cells: a potential mechanism for position-effect variegation. *Proc. Natl. Acad. Sci.* 2003b; 100:1820-5.
- Nickel BE, Davie JR. Structure of polyubiquitinated histone H2A. *Biochemistry* 1989;28:964-68.
- Nickel BE, Allis CD, Davie JR. Ubiquitinated histone H2B is preferentially located in transcriptionally active chromatin. *Biochemistry* 1989;28:958-963.
- Nielsen PR, Nietlispach D, Mott HR, Callaghan J, Bannister A, Kouzarides T, Murzin AG, Murzina NV, Laue ED. Structure of the HP1 chromodomain bound to histone H3 methylated at lysine 9. *Nature* 2002; 416:103-7.
- Nielsen SJ, Schneider R, Bauer UM, Bannister AJ, Morrison A, O'Carroll D, Firestein R, Cleary M, Jenuwein T, Herrera RE, Kouzarides T. Rb targets histone H3 methylation and HP1 to promoters. *Nature* 2001;412:561-5.
- Nishida T, Tanaka H, Yasuda H. A novel mammalian Smt3-specific isopeptidase 1 (SMT3IP1) localized in the nucleolus at interphase. *Eur. J. Biochem.* 2000;267:6423-7.

- North BJ, Marshall BL, Borra MT, Denu JM, Verdin E. The human Sir2 ortholog SIRT2, is an NAD<sup>+</sup>-dependent tubulin deacetylase. *Mol. Cell* 2003;437-44.
- North BJ, Verdin E. Sirtuins: Sir2-related NAD-dependent protein decetylases. *Genome Biol.* 2004;5:224.
- Norton VG, Imai BS, Yau P, Bradbury EM. Histone acetylation reduces nucleosome core particle linking number change. *Cell* 1989;57:449-57.
- Nowak SJ, Corces VG. Phosphorylation of histone H3 correlates with transcriptionally active loci. *Genes Dev.* 2000;14:3003-13.
- OMS *Obesidad y sobrepeso*. Organización Mundial de la Salud. Disponible en: <http://www.who.int/mediacentre/factsheets/fs311/es/index.html>. Accesado el 4 de junio 2008.
- O'Carrol D, Scherthan H, Peter AH, Opravil S, Haynes AR, Laible G, Rea S, Schmid M, Lebersorger A, Jerratsch M, Sattler L, Mattei MG, Denny P, Brown SD, Schweizer D, Jenuwein T. Isolation and characterization of Suv39h2, a second histone H3 methyltransferase gene that displays testis-specific expression. *Mol. Cell Biol.* 2000;20:9423-33.
- Ohara K, Xu HG, Mori N, Suzuki Y, Xu DS, Wang ZC. Anticipation and imprinting in schizophrenia. *Biol. Psychiatry*. 1997;42:760-66.
- Okamoto I, Otte AP, Allis CD, Reinberg D, Heard E. Epigenetic dynamics of imprinted X inactivation during mouse development. *Science* 2004;303:644-9.
- Okano M, Xie S, Li E. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nature Genet.* 1998a;19:219-20.
- Okazaki IJ, Zolkiewska A, Takada T, Moss J. Characterization of mammalian ADP-ribosylation cycles. *Biochimie* 1995; 77:319-25.
- Okazaki IJ, Moss J. Mono-ADP-ribosylation: a reversible posttranslational modification of proteins. *Adv. Pharmacol.* 1996; 35:247-80.
- Okazaki IJ, Moss J. Characterizacion of glycosylphosphatidyl-inositol-anchored, secreted, and intracellular vertebrate mono-ADP-ribosyltransferases. *Annu. Rev. Nutr.* 1999; 19:485-509.
- Okazaki R, Okazaki T, Sakabe K. Mechanism of DNA replication possible discontinuity of DNA chain growth. *Jpn. J. Med. Sci. Biol.* 1967; 20:255-60.
- Okoji RS, Yu RC, Maronpot RR, Froines JR. Sodium arsenite administration via drinking water increases genome-wide and Ha-ras DNA hypomethylation in methyl-deficient C57BL/6J mice. *Carcinogenesis* 2002;23:777-85.
- Olaharski AJ, Rine J, Marshall BL, Babiarz J, Zhang L, Verdin E, Smith MT. The flavouring agent dihydrocoumarin reverses epigenetic silencing and inhibits sirtuina deacetylases. *PLoS Genet.* 2005;1:e77.
- Oommen AM, Griffin JB, Sarta G, Zempleni J. Roles for nutrients in epigenetic events. *J. Nutr. Biochem.* 2005; 16:74-7.
- Ord MG, Stocken LA. Adenosine diphosphate ribosylated histones. *Biochem. J.* 1977;161:583-92.
- Ordoval JM. HDL genetics: candidate genes, genome wide scans and gene-environment interactions. *Cardiovasc. Drugs Ther.* 2002;16:273-81.
- Oster H. Sex-based differences in gene transmisión and gene expresión. *Lupus* 1999;8:365-9.
- Ottman R. Gene-environment interaction; definitions and study designs. *Prev. Med.* 1996;25:764-70.
- Otto H, Reche PA, Bazan F, Dittmar K, Haag F, Koch-Nolte F. In silico characterization of the family of PARP-like poly(ADP-ribosyl)transferases (pARTs). *BMC Genomics* 2005;6:139.
- Ou JN, Torrisani J, Unterberger A, Provençal N, Shikimi K, Karimi M, Ekström TJ, Szyf M. Histone deacetylase inhibitor trichostatin A induces global and gene-specific DNA demethylation in human cancer cell lines. *Biochem. Pharmacol.* 2007;73:1297-307.
- Owen DJ, Ornaghi P, Yang JC, Lowe N, Evans PR, Ballario P, Neuhaus D, Filetici P, Travers AA. The structural basis for the recognition of acetylated histone H4 by the bromodomain of histone acetyltransferase Gcn5p. *EMBO J.* 2000;19:6141-9.

- Paik WK, Kim S. Enzymatic methylation of protein fraction from calf thymus nuclei. *Biochem. Biophys. Res. Commun.* 1967;29:14-20.
- Painter RC, Roseboom TJ, Bleker OP. Prenatal exposure to the Dutch famine and disease in later life: an overview. *Repro. Toxicol.* 2005; 20:345-52.
- Pal S, Vishwanath SN, Erdjument-Bromage H, Tempst P, Sif S. Human SWI/SNF-associated PRMT5 methylates histone H3 arginine 8 and negatively regulates expression of ST7 and NM23 tumor suppressor genes. *Mol. Cell Biol.* 2004;24:9630-45.
- Pandey SC, Ugale R, Zhang H, Tang L, Prakash A. Brain chromatin remodeling: a novel mechanism of alcoholism. *J. Neurosci.* 2008;28:3729-37.
- Panning B, Jaenisch R. DNA hypomethylation can activate Xist expression and silence X-linked genes. *Genes Dev.* 1996;10:1991-2002.
- Parlow MH, Haas AL, Lough J. Enrichment of ubiquitinated histone H2A in a low salt extract of micrococcal nuclease-digested myotube nuclei. *J. Biol. Chem.* 1990;265:7507-12.
- Pauer LR, Olivares J, Cunningham C, Williams A, Grove W, Kraker A, Olson S, Nemunaitis J. Phase I study of oral CI-994 in combination with carboplatin and paclitaxel in the treatment of patients with advanced solid tumors. *Cancer Invest.* 2004;22:886-96.
- Pauling L. Orthomolecular psychiatry. Varying the concentrations of substances normally present in the human body may control mental disease. *Science* 1968;160:265-71.
- Pazin MJ, Kadonaga JT. What's up and down with histone deacetylation and transcription? *Cell* 1997;89:325-8.
- Peart MJ, Smyth GK, van Laar RK, Bowtell DD, Richon VM, Marks PA, Holloway AJ, Johnstone RW. Identification and functional significance of genes regulated by structurally different histone deacetylase inhibitors. *Proc. Natl. Acad. Sci. USA* 2005;102:3697-702
- Peaston AE, Whitelaw E. Epigenetics and phenotypic variation in mammals. *Mamm. Genome* 2006; 17:365-74.
- Pei Y, Niu L, Lu F, Liu C, Zhai J, Kong X, Cao X. Mutations in type II protein arginine methyltransferase AtPRMT5 result in pleiotropic developmental defects in arabidopsis. *Plant Physiol.* 2007;144:1913-23.
- Pembrey ME. Time to take epigenetic inheritance seriously. *Eur. J. Hum. Genet.* 2002; 10:669-71.
- Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K, Sjöström M, Golding J. Sex-specific, male-line transgenerational responses in humans. *Eur. J. Hum. Genet.* 2006;14:159-66.
- Peña P, Davrazou F, Shi X, Walter KL, Verkhusha VV, Gozani O, Zhao R, Kutateladze TG. Molecular mechanism of histone H3K4me3 recognition by plant homeodomain of ING2. *Nature* 2006;442:100-3.
- Perez EA, Pusztai L, van de Vijver M. Improving patient care through molecular diagnostics. *Semin. Oncol.* 2004;31:14-20.
- Perez-Lamigueiro MA, Alvarez-Gonzalez R. Polynucleosomal synthesis of poly(ADP-ribose) causes chromatin unfolding as determined by micrococcal nuclease digestion. *Ann. NY Acad. Sci.* 2004;1030:593-8.
- Perrod S, Cockell MM, Laroche T, Renauld H, Ducrest AL, Bonnard C, Gasser SM. A cytosolic NAD-dependent deacetylase, Hst2p, can modulate nucleolar and telomeric silencing in yeast. *EMBO J.* 2001;20:197-209.
- Peters AH, O'Carrol D, Scherthan H, Mechtler K, Sauer S, Schöfer C, Weipoltshammer K, Pagani M, Lachner M, Kohlmaier A, Opravil S, Doyle M, Sibilia M, Jenuwein T. Loss of Suv39h histone methyltransferases impairs mammalian heterochromatin and genome stability. *Cell* 2001;107:323-37.
- Peters AH, Kubicek S, Mechtler K, O'Sullivan RJ, Derijck AA, Perez-Burgos L, Kohlmaier A, Opravil S, Tachibana M, Shinkai Y, Martens JH, Jenuwein T. Partitioning and plasticity of repressive histone methylation states in mammalian chromatin. *Mol. Cell* 2003;12:1577-89.
- Peters DM, Griffin B, Stanley JS, Beck MM, Zempleni J. Exposure to UV light causes increased biotinylation of histones in Jurkat cells. *Am. J. Physiol. Cell Physiol.* 2002;283:C878-84,
- Peters JM. SCF and APC: the Yin and Yang of cell cycle regulated proteolysis. *Curr. Opin. Cell Biol.* 1998;10:759-68.

- Petrelli F, Marsili G, Moretti P. RNA, DNA, histones and interactions between histone proteins and DNA in the liver of biotin-deficient rats. *Biochem. Exp. Biol.* 1976;12:461-5.
- Petrelli F, Coderoni S, Moretti P, Paparelli M. Effect of biotin on phosphorylation, acetylation, methylation of rat liver histones. *Mol. Biol. Rep.* 1978;4:87-92.
- Petronis A. Human morbid genetics revisited: relevance of epigenetics. *Trends Genet.* 2001;17:142-6.
- Petruck S, Sedkov Y, Smith S, Tillib S, Kraevski V, Nakamura T, Canaani E, Croce CM, Mazo A. Trithorax and dCBP acting in a complex to maintain expression of a homeotic gene. *Science* 2001;294:1331-4.
- Pflum MK, Tong JK, Lance WS, Schreiber SL. Histone deacetylase 1 phosphorylation promotes enzymatic activity and complex formation. *J. Biol. Chem.* 2001;276:47496-507.
- Pham AD, Sauer F. Ubiquitin-activating/conjugating activity of TAF<sub>II</sub>250, a mediator of activation of gene expression in *Drosophila*. *Science* 2000;289:2357-60.
- Picard F, Kurtev M, Chung N, Topark-Ngarm A, Senawong T, Machado de Oliveira R, Leid M, McBurney MW, Guarente L. Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR-g. *Nature* 2005;434:113-8.
- Pichler A, Gast A, Seeler JS, Dejean A, Melchior F. The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* 2002;108:109-20.
- Pickart CM. Mechanisms underlying ubiquitination. *Annu. Rev. Biochem.* 2001;70:503-33.
- Piekorz RL, Robey R, Sandor V, Bakke S, Wilson WH, Dahmoush L, Kingma DM, Turner ML, Altermus R, Bates SE. Inhibitor of histone deacetylation, depsipeptide (FR901228), in the treatment of peripheral and cutaneous T-cell lymphoma: a case report. *Blood* 2001;98:2865-8.
- Pispa J. Animal biotinidase. *Ann. Med. Exp. Biol. Fenn.* 1965;43:4-39.
- Plath K, Fang J, Mlynarczyk-Evans SK, Cao R, Worringer KA, Wang H, de la Cruz CC, Otte AP, Panning B, Zhang Y. Role of histone H3 lysine 27 methylation in X inactivation. *Science* 2003;300:131-5.
- Pogo BGT, Allfrey VG, Mirsky AE. RNA synthesis and histone acetylation during the course of gene activation in lymphocytes. *Proc. Natl. Acad. Sci.* 1966;55:805-12.
- Pogribny IP, Ross SA, Tryndyak VP, Pogribna M, Poirier LA, Karpinets TV. Histone H3 lysine 9 and H4 lysine 20 trimethylation and the expression of Suv4-20h2 and Suv-39h1 histone methyltransferases in hepatocarcinogenesis induced by methyl deficiency in rats. *Carcinogenesis* 2006;27:1180-6.
- Poirier GG, de Murcia G, Jongstra-Bilen J, Niedergang C, Mandal P. Poly(ADP-ribosyl)ation of polynucleosomes causes relaxation of chromatin structure. *Proc. Natl. Acad. Sci. USA* 1982;79:3423-7.
- Poirier LA. The role of methionine in carcinogenesis in vivo. *Adv. Exp. Med. Biol.* 1986;206:269-82.
- Poirier LA, Brown AT, Fink LM, Wise CK, Randolph CJ, Delongchamp RR, Fonseca VA, Blood S. Adenosylmethionine concentrations and lymphocyte methylenetetrahydrofolate reductase activity in diabetes mellitus and diabetic nephropathy. *Metabolism* 2001;50:1014-8.
- Poirier LA, Vlasova TI. The prospective role of abnormal methyl metabolism in cadmium toxicity. *Environ. Health Perspect.* 2002; 110:793-5.
- Pompeia C, Hodge DR, Plass C, Wu YZ, Marquez VE, Kelley JA, Farrar WL. Microarray analysis of epigenetic silencing of gene expression in the KAS-6/1 multiple myeloma cell line. *Cancer Res.* 2004;64:3465-73.
- Portoso M, Cavalli G. The role of RNAi and noncoding RNAs in polycomb mediated control of gene expression and genomic programming. En: *RNA and the Regulation of Gene Expression: a Hidden Layer of Complexity*. K.V. Morris, Caister Academic Press; 2008.
- Posfai J, Bhagwat AS, Posfai G, Roberts RJ. Predictive motifs derived from cytosine methyltransferases. *Nucleic Acids Res.* 1989;17:2421-35.
- Post WS, Goldschmidt-Clermont PJ, Wilhede CC, Heldman AW, Sussman MS, Ouyang P, Milliken EE, Issa JP. Methylation of the estrogen receptor gene is associated with aging and atherosclerosis in the cardiovascular system. *Cardiovasc. Res.* 1999;43:985-91.
- Potter GB, Beaudoin GMJ III, DeRenzo CL, Zarach JM, Chen SH, Thompson CC. The hairless gene mutated in congenital hair loss disorders encodes a novel nuclear receptor corepressor. *Genes Dev.* 2001;15:2687-701.

- Pradhan S, Kim GD. The retinoblastoma gene product interacts with maintenance human DNA (cytosine-5) methyltransferase and modulates its activity. *EMBO J.* 2002;21:779-88.
- Prakash S, Foster BJ, Meyer M, Wozniak A, Heilbrun LK, Flaherty L, Zalupski M, Radulovic L, Valdivieso M, LoRusso PM. Chronic oral administration of CI-994: a phase 1 study. *Invest. New Drugs* 2001;19:1-11.
- Pray-Grant MG, Schieltz D, McMahon SJ, Wood JM, Kennedy EL, Cook RG, Workman JL, Yates JR, Grant PA. The novel SILK histone acetyltransferase complex functions in the yeast retrograde response pathway. *Mol. Cell Biol.* 2002;22:8774-86.
- Pray-Grant MG, Daniel JA, Schieltz D, Yates JR, Grant PA. Chd1 chromodomain links histone H3 methylation with SAGA- and SLIK-dependent acetylation. *Nature* 2005;433:434-8.
- Ptak C, Petronis A. Epigenetics and complex disease: from etiology to new therapeutics. *Annu. Rev. Pharmacol. Toxicol.* 2008; 48:257-76.
- Qiu C, Sawada K, Zhang X, Cheng X. The PWWP domain of mammalian DNA methyltransferase Dnmt3b defines a new family of DNA-binding folds. *Nat. Struct. Biol.* 2002;9:217-24.
- Qiu J. Epigenetics: unfinished symphony. *Nature* 2006; 441:143-45.
- Quddus J, Jonson KJ, Gavalchin J, Amento EP, Chriss CE, Yung RL, Richardson BC. Treating activated CD4+ T cells with either of two distinct DNA methyltransferase inhibitors, 5-azacytidine or procainamide, is sufficient to cause a lupus-like disease in syngeneic mice. *J. Clin. Invest.* 1993;92:38-53.
- Ran Q, Pereira-Smith OM. Identification of an alternatively spliced form of the Tat interactive protein. (Tip60), Tip60beta. *Gene* 2000;258:141-6.
- Rauch T, Li H, Wu X, Pfeifer GP. MIRA-assisted microarray analysis, a new technology for the determination of DNA methylation patterns: identifies frequent methylation of homeodomain-containing genes in lung cancer cells. *Cancer Res.* 2006;66:7939-47.
- Raysam GV, Wendling O, Angrand PO, Mark M, Niederreither K, Song L, Lerouge T, Hager GL, Chambon P, Losson R. NSD1 is essential for early post-implantation development and has a catalytically active SET domain. *EMBO J.* 2003;22:3153-63.
- Rea S, Eisenhaber F, O'Carroll D, Strahl BD, Sun ZW, Schmid M, Opravil S, Mechtler K, Ponting CP, Allis CD, Jenuwein T. Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* 2000;406:593-99.
- Reifsnyder C, Lowell J, Clarke A, Pillus L. Yeast SAS silencing genes and human genes associated with AML and HIV-1 Tat interactions are homologous with acetyltransferases. *Nature Genet.* 1996;14:42-9.
- Reik W, Walter J. Genomic imprinting: parental influence on the genome. *Nat. Rev. Genet.* 2001;2:21-32.
- Reiss Y, Heller H, Hershko A. Binding sites of ubiquitina-protein ligase. Binding of ubiquitina-protein conjugates and of ubiquitin-carrier protein. *J. Biol. Chem.* 1989;264:10378-83.
- Remiszewski SW, Sambucetti LC, Bair KW, Bontempo J, Cesarz D, Chandramouli N, Chen R, Cheung M, Cornell-Kennon S, Dean K, Diamantidis G, France D, Green MA, Howell KL, Kashi R, Kwon P, Lassota P, Martin MS, Mou Y, Perez LB, Sharma S, Smith T, Sorensen E, Taplin F, Trogani N, Versace R, Walker H, Weltchek-Engler S, Wood A, Wu A, Atadja P. N-hydroxy-3-phenyl-2-propenamides as novel inhibitors of human disease deacetylase with in vivo antitumor activity: discovery of (2E)-N-hydroxy-3-[4-[(2-hydroxyethyl)[2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]-2-propenamide (NYP-LAO84). *J. Med. Chem.* 2003;46:4609-24.
- Rezai-Zadeh N, Zhang X, Namour F, Fejer G, Wen YD, Yao YL, Gyory I, Wright K, Seto E. Targeted recruitment of a histone H4-specific methyltransferase by the transcription factor YY1. *Genes Dev.* 2003;17:1019-29.
- Rice JC, Briggs SD, Ueberheide B, Barber CM, Shabanowitz J, Hunt DF, Shinkai Y, Allis CD. Histone methyltransferase direct different degrees of methylation to define distinct chromatin domains. *Mol. Cell* 2003;12:1591-8.
- Richardson B. DNA methylation and autoimmune disease. *Clin. Immunol.* 2003; 109:72-9.
- Richardson B, Scheinbart L, Strahler J, Gross L, Hanash S, Johnson M. Evidence for impaired T cell DNA methylation in systemic lupus erythematosus and rheumatoid arthritis. *Arthritis Rheum.* 1990; 33:1665-73.

- Richardson BC. Role of DNA methylation in the regulation of cell function: autoimmunity, aging and cancer. *J. Nutr.* 2002;132:2401S-2405S.
- Richon VM, Webb Y, Merger R, Sheppard T, Jursic B, Ngo L, Civoli F, Breslow R, Rifkind RA, Marks PA. Second generation hybrid polar compounds are potent inducers of transformed cell differentiation. *Proc. Natl. Acad. Sci. USA* 1996;93:5705-8.
- Riggs AD. X inactivation, differentiation, and DNA methylation. *Cytogenet. Cell Genet.* 1975; 15:9-25.
- Riggs AD, Xiong Z, Wang L, LeBon JM. Methylation dynamics, epigenetic fidelity and X chromosome structure. *Novartis Found Symp.* 1998;214:214-25; discussion 225-32.
- Ringrose L, Paro R. Epigenetic regulation of cellular memory by the polycomb and trithorax group proteins. *Annu. Rev. Genet.* 2004;38:413-43.
- Rippmann JF, Damm K, Schnapp A. Functional characterization of the poly(ADP-ribose) polymerase activity of tankyrase 1, a potential regulator of telomer length. *J. Mol. Biol.* 2002;323:217-24.
- Robertson KD. DNA methylation, methyltransferases, and cancer. *Oncogene* 2001;20:3139-55.
- Robertson KD. DNA methylation and chromatin-unraveling the tangled web. *Oncogene* 2002;21:5361-79.
- Robertson KD. DNA methylation and human disease. *Nat. Rev. Genet.* 2005;6:597-610.
- Robertson KD, Wolffe AP. DNA methylation in health and disease. *Nature Rev. Genet.* 2000;1:11-9.
- Robertson KD, Uzvolgyi E, Liang G, Talmadge C, Sumegi J, Gonzales FA, Jones PA. The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Res.* 1999;27:2291-8.
- Robertson KD, Ait-Si-Ali S, Yokochi T, Wade PA, Jones PL, Wolffe AP. DNMT1 forms a complex with Rb, E2F1 and HDAC1 and represses transcription from E2F-responsive promoters. *Nat. Genet.* 2000a;25:338-42.
- Robertson KD, Keyomarsi K, Gonzales FA, Velicescu M, Jones PA. Differential mRNA expression of the human DNA methyltransferases (DNMTs) 1, 3a and 3b during the G(0)/G(1) to S phase transition in normal and tumor cells. *Nucleic Acids Res.* 2000b; 28:2108-13.
- Robzyk K, Recht J, Osley MA. Rad6-dependent ubiquitination of histone H2B in yeast. *Science* 2000;287:501-4.
- Rodgers JT, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1 $\alpha$  and SIRT1. *Nature* 2005;434:113-8.
- Rogakou EP, Boon C, Redon C, Bonner WM. Megabase chromatin domains involved in DNA double-strand breaks *in vivo*. *J. Cell Biol.* 1999;146:905-16.
- Roguev A, Schaft D, Shevchenko A, Aasland R, Shevchenko A, Stewart AF. High conservation of the Set1/Rad6 axis of histone 3 lysine 4 methylation in budding and fission yeasts. *J. Biol. Chem.* 2003;278:8487-93.
- Rollins RA, Haghghi F, Edwards JR, Das R, Zhang MQ, Ju J, Bestor TH. Large-scale structure of genomic methylation patterns. *Genome Res.* 2006;16:157-63.
- Rosato RR, Almenara JA, Dai Y, Grant S. Simultaneous activation of the intrinsic and extrinsic pathways by histone deacetylase (HDAC) inhibitors and tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) synergistically induces mitochondrial damage and apoptosis in human leukemia cells. *Mol. Cancer Ther.* 2003;2:1273-84.
- Ross SA Diet and DNA methylation interactions in cancer prevention. *Ann. NY Acad. Sci.* 2003;983:197-207.
- Ross S, Best JL, Zon LI, Gill G. SUMO-1 modification represses Sp3 transcriptional activation and modulates its subnuclear localization. *Mol. Cell* 2002;10:831-42.
- Rossmann RS. Comparison of super-secondary structures in proteins. *J. Mol. Biol.* 1973;76:241-56.
- Roth SY, Denu JM, Allis CD. Histone acetyltransferases. *Annu. Rev. Biochem.* 2001;70:81-120.
- Rountree MR, Bachman KE, Baylin SB. DNMT1 binds HDAC2 and a novel corepressor DMAP1, to form a complex at replication foci. *Nat. Genet.* 2000;25:269-77.
- Rowling MJ, McMullen MH, Schalinske KL. Vitamin A and its derivatives induce hepatic glycine N-methyltransferase and hypomethylation of DNA in rats. *J. Nutr.* 2002;132:365-9.

- Ruefli AA, Ausserlechner MJ, Bernhard D, Sutton VR, Tainton KM, Kofler R, Smyth MJ, Johnstone RW. The histone deacetylase inhibitor and chemotherapeutic agent suberoylanilide hydroxamic acid (SAHA) induces a cell-death pathway characterized by cleavage of Bid and production of reactive oxygen species. *Proc. Natl. Acad. Sci. USA* 2001;98:10833-8.
- Ruijter AJM de, van Gennip AH, Caron HN, Kemp S, van Kuilenburg ABP. Histone deacetylases (HDACs): characterization of the classical HDAC family. *Biochem. J.* 2003;370:737-49.
- Ryan RF, Schultz DC, Ayyanathan K, Singh PB, Friedman JR, Fredericks WJ, Rauscher FJ. KAP-1 corepressor protein interacts and colocalizes with heterochromatic and euchromatic HP1 proteins: a potential role for Kruppel-associated box-zinc finger proteins in heterochromatin-mediated gene silencing. *Mol. Cell Biol.* 1999;19:4366-78.
- Ryan QC, Headlee D, Acharya M, Sparreboom A, Trepel JB, Ye J, Figg WD, Hwang K, Chung EJ, Murgo A, Melillo G, Elsayed Y, Monga M, Kalnitskiy M, Zwiebel J, Sausville EA. Phase I and pharmacokinetic study of MS-275, a histone deacetylase inhibitor, in patients with advanced refractory solid tumors or lymphoma. *J. Clin. Oncol.* 2005;23:3912-22.
- Rybicki E. PCR primer design and reaction optimisation. Molecular Biology Techniques Manual. Disponible en: <http://www.mcb.uct.ac.za/pcoptim.htm> Accesado el 1 de junio 2008.
- Sachdev S, Bruhn L, Sieber H, Pichler A, Melchior F, Grosschedl R. PIASy, a nuclear matrix-associated SUMO E3 ligase, represses LEF1 activity by sequestration into nuclear bodies. *Genes Dev.* 2001;15:3088-103.
- Saikawa Y, Kubota T, Maeda S, Otani Y, Kumai K, Kitajima M. Inhibition of DNA methyltransferase by antisense oligodeoxynucleotide modifies cell characteristics in gastric cancer cell lines. *Oncol. Rep.* 2004;12:527-31.
- Saladrigas MV, Claros MG, González D. Vocabulario de inglés-español de bioquímica y biología molecular. Panace@. Disponible en: <http://www.biorom.uma.es/contenido/Glosario/>. Acessado en 30 abril 2008.
- Saleh A, Schieltz D, Ting N, McMahon SB, Litchfield DW, Yates JR 3rd, Lees-Miller SP, Cole MD, Brandl CJ. Tra1p is a component of the yeast Ada.Spt transcriptional regulatory complexes. *J. Biol. Chem.* 1998;273:26559-65.
- Sallmann FR, Vodenicharov MD, Wang ZQ, Poirier GG. Characterization of sPARP-1. An alternative product of PARP-1 gene with poly(ADP-ribose) polymerase activity independent of DNA strand breaks. *J. Biol. Chem.* 2000;275:15504-11.
- Samaco RC, Hogart A, LaSalle JM. Epigenetic overlap in autism-spectrum neurodevelopmental disorders: MECP2 deficiency causes reduced expression of UBE3A and GABRB3. *Hum. Mol. Genet.* 2005;14:483-92.
- Sánchez-Céspedes M, Esteller M, Wu L, Nawroz-Danish H, Yoo GH, Koch WM, Jen J, Herman JG, Sidransky D. Gene promoter hypermethylation in tumors and serum of head and neck cancer patients. *Cancer Res.* 2000;60:892-5.
- Sanders SL, Portoso M, Mata J, Bahler J, Allshire RC, Kouzarides T. Methylation of histone H4 lysine 20 controls recruitment of Crb2 to sites of DNA damage. *Cell* 2004;119:603-14.
- Sandor V, Bakke S, Robey RW, Kang MH, Blagosklonny MV, Bender J, Brooks R, Piekarz RL, Tucker E, Figg WD, Chan KK, Goldspiel B, Fojo AT, Balcerzak SP, Bates SE. Phase I trial of the histone deacetylase inhibitor, depsipeptide (FR901228, NSC 630176), in patients with refractory neoplasms. *Clin. Cancer Res.* 2002;8:718-28.
- Santi DV, Garrett CE, Barr PJ. On the mechanism of inhibition of DNA-cytosine methyltransferases by cytosine analogs. *Cell* 1983;33:9-10.
- Santini V, Kantarjian HM, Issa JP. Changes in DNA methylation and neoplasia: pathophysiology and therapeutic implications. *Ann. Intern. Med.* 2001;134:573-86.
- Santos-Rosa H, Caldas C. Chromatin modifier enzymes, the histone code and cancer. *Eur. J. Cancer.* 2005; 41:2381-2402.
- Sassone-Corsi P, Mizzen CA, Cheung P, Crosio C, Monaco L, Jacquot S, Hanauer A, Allis CD. Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science* 1999;285:886-91.

- Sayito H, Hinche J. Functional heterogeneity of small ubiquitin-related protein modifiers SUMO-1 versus SUMO-2/3. *J. Biol. Chem.* 2000;275:6252-8.
- Scarano MI, Strazzullo M, Matarazzo MR, D'Esposito M. DNA methylation 40 years later: its role in human health and disease. *J. Cell Physiol.* 2005;204:21-35.
- Schaft D, Rogev A, Kotovic KM, Shevchenko A, Sarov M, Shevchenko A, Neugebauer KM, Stewart AF. The histone 3 lysine 36 methyltransferase, SET2, is involved in transcriptional elongation. *Nucleic Acids Res.* 2003;31:2475-82.
- Schanen NC. Epigenetics of autism spectrum disorders. *Hum. Mol. Genet.* 2006;15:R138-R150.
- Scheerger SB, Zempleni J. Expression of oncogenes depends on biotin in human small cell lung cancer cells NCI-H69. *Int. J. Vitam. Nutr. Res.* 2003;73:461-7.
- Scheffner M, Nuber U, Huibregtse JM. Protein ubiquitination involving an E1-E2-E3 enzyme ubiquitin thioester cascade. *Nature* 1995;373:81-3.
- Scheinbart LS, Jonson MA, Gross LA, Eldelstein SR, Richardson BC. Procainamide inhibits DNA methyltransferase in a human T cell line. *J. Rheum.* 1991;18:530-40.
- Schindler U, Beckmann H, Cashmore AR. HAT3.1, a novel *Arabidopsis* homeodomain protein containing a conserved cysteine-rich region [abstract]. *Plant J.* 1993; 4:137-50. <http://www.ncbi.nlm.nih.gov/pubmed/8106082>. Accesado el 16 de mayo 2008.
- Schneider R, Bannister AJ, Kouzarides T. Unsafe SETs: histone lysine transferases and cancer. *Trends Biochem. Sci.* 2002;27:396-402.
- Schoenherr CJ, Levorse JM, Tilghman SM. CTCF maintains differential methylation at the Igf2/H19 locus. *Nature Genet.* 2003;2003;66-9.
- Schotta H, Lachner M, Sarma K, Ebert A, Sengupta R, Reuter G, Reinberg D, Jenuwein T. A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev.* 2004;18:1251-62.
- Schreiber V, Ame JC, Dolle P, Schultz I, Rinaldi B, Fraulob V, Méniére-de Murcia J, de Murcia G. Poly(ADP-ribose) polymerase-2 (PARP-2) is required for efficient base excision DNA repair in association with PARP-1 and XRCC1. *J. Biol. Chem.* 2002;277:23028-36.
- Schroer RJ, Phelan MC, Michaelis RC, Crawford EC, Skinner SA, Cuccaro M, Simensen RJ, Bishop J, Skinner C, Fender D, Stevenson RE. Autism and maternally derived aberrations of chromosome 15q. *Am. J. Med. Genet.* 1998;76:327-36.
- Schulze TG, Chen YS, Badner JA, McInnis MG, DePaulo JRJ, McMahon FJ. Additional, physically ordered markers increase linkage signal for bipolar disorder on chromosome 18q22. *Biol. Psychiatry* 2003;53:239-43.
- Schurter BT, Joh SS, Chen D, Bunick GJ, Harp JM, Hanson BL, Henschen-Edman A, Mackay DR, Stallcup MR, Aswad DW. Methylation of histone H3 by coactivator-associated arginine methyltransferase 1. *Biochem.* 2001;40:5747-56.
- Scott JM, Weir DG. Folic acid, homocysteine and one-carbon metabolism: a review of the essential biochemistry. *J. Cardiovas. Risk* 1998;5:223-227.
- Sealy L, Chalkley R. DNA associated with hyperacetylated histone is preferentially by DNase I. *Nucleic Acids Res.* 1978;5:1863-76.
- Sedkov Y, Cho E, Petruk S, Cherbas L, Smith ST, Jones RS, Cherbas P, Canaani E, Jaynes JB, Mazo A. Methylation at lysine 4 of histone H3 in ecdysone-dependent development of *Drosophila*. *Nature* 2003;426:78-83.
- Seigneurin-Berny D, Verdel A, Curtelet S, Lemercier C, Garin J, Rousseaux S, Khochbin S. Identification of components of the murine histone deacetylase 6 complex: link between acetylation and ubiquitination signalling pathways. *Mol. Cell Biol.* 2001;21:8035-44.
- Seman M, Adriouch S, Haag F, Koch-Nolte F. Ecto-ADP-ribosyltransferases (ARTs): emerging actors in cell communication and signalling. *Curr. Med. Chem.* 2004;11:857-72.
- Sengupta N, Seto E. Regulation of histone deacetylase activities. *J. Cell Biochem.* 2004;93:57-67.

- Sgarra R, Diana F, Bellarosa C, Dekleva V, Rustighi A, Toller M, Manioletti G, Giancotti V. During apoptosis of tumor cells HMGA1a protein undergoes methylation: identification of the modification site by mass spectrometry. *Biochemistry* 2003;42:3575-85.
- Shen L, Waterland RA. Methods of DNA methylation analysis. *Curr. Opin. Clin. Nutr. Metab. Care* 2007; 10:576-81.
- Shen L, Guo Y, Chen X, Ahmed S, Issa JP. Optimizing annealing temperature overcomes bias in bisulfite PCR methylation analysis. *Biotechniques* 2007; 42:50-2.
- Shi H, Maier S, Nimmrich I, Yan PS, Caldwell CW, Olek A, Huang TH. Oligonucleotide-based microarray for DNA methylation analysis: principles and applications. *J. Cell Biochem.* 2003;88:138-43.
- Shi X, Hong T, Walter KL, Ewalt M, Michishita E, Hung T, Carney D, Peña P, Lan F, Kaadige MR, Lacoste N, Cayrou C, Davrazou F, Saha A, Cairns BR, Ayer DE, Kutateladze TG, Shi Y, Côté J, Chua KF, Gozani O. ING2 PHD domain links histone H3 lysine 4 methylation to active gene repression. *Nature* 2006;442:96-9.
- Shi Y, Lan F, Matson C, Mulligan P, Whetstone JR, Cole PA, Casero RA, Shi Y. Histone demethylation mediated by the nuclear amine oxydase homolog LSD1. *Cell* 2004;119:941-53.
- Shi YJ, Matson C, Lan F, Iwase S, Bata T, Shi Y. Regulation of LSD1 histone demethylase activity by its associated factors. *Mol. Cell* 2005;19:857-64.
- Shioi Y, Eisenman RN. Histone sumoylation is associated with transcriptional repression. *Proc. Natl. Acad. Sci. USA* 2003;100:13225-30.
- Sibley CP, Coan PM, Ferguson-Smith AC, Dean W, Hughes J, Smith P, Reik W, Burton GJ, Fowden AL, Constâncio M. Placental-specific insulin-like growth factor 2 (Igf2) regulates the diffusional exchange characteristics of the mouse placenta. *Proc. Natl. Acad. Sci. USA* 2004;101:8204-8.
- Siedlecki P, Zielenkiewicz P. Mammalian DNA methyltransferases. *Acta. Biochim. Pol.* 2006;53:245-56.
- Siedlecki P, Boy RH, Comagic S, Schirrmacher R, Wiessler M, Zielenkiewicz P, Suhai S, Lyko F. Establishment and functional validation of a structural homology model for human DNA methyltransferase 1. *Biochem. Biophys. Res. Commun.* 2003;306:558-63.
- Silva JM, Domínguez G, García JM, González R, Villanueva MJ, Navarro F, Provencio M, San Martín S, España P, Bonilla F. Presence of tumor DNA in plasma of breast cancer patients: clinicopathological correlations. *Cancer Res.* 1999;59:3251-6.
- Silva J, Mark W, Zvetkova I. Establishment of histone H3 methylation on the inactive X chromosome requires transient recruitment of Eed-Enx1 polycomb group complexes. *Dev. Cell* 2003;4:481-95.
- Singh SM, Murphy B, O'Reilly RL. Involvement of gene-diet/drug interaction in the DNA methylation and its contribution to complex diseases: from cancer to schizophrenia. *Clin. Genet.* 2003;64:451-60.
- Skalníková M, Bártová E, Ulman V, Matula P, Svoboda D, Harnicarová A, Kozubek M, Kozubek S. Distinct patterns of histone methylation and acetylation in human interphase nuclei. *Physiol. Res.* 2007;56:797-806.
- Smale ST, Kadonaga JT. The RNA polymerase II core promoter. *Annu. Rev. Biochem.* 2003; 72:449-79.
- Smit AF, Riggs AD. Triggers and other DNA transposons fossils in the human genome. *Proc. Natl. Acad. Sci. USA* 1996;13:335-40.
- Smith ER, Belote JM, Schiltz RL, Yang XJ, Moore PA, Berger SL, Nakatani Y, Allis CD. Cloning of *Drosophila* GCN5: conserved features among metazoan GCN5 family members. *Nucleic Acids Res* 1998a; 26:2948-54.
- Smith ER, Eisen A, Gu W, Sattah M, Pannuti A, Zhou J, Cook RG, Lucchesi JC, Allis CD. ESA1 is a histone acetyltransferase that is essential for growth in yeast. *Proc. Natl. Acad. Sci. USA* 1998b; 95:3561-5.
- Smith JA, Stocken LA. Chemical and metabolic properties of adenosine diphosphate ribose derivatives of nuclear proteins. *Biochem. J.* 1975; 147:523-9.
- Smith KP, Benjamín RC, Moss J, Jacobson MK. Identification of enzymatic activities which process protein bound mono(ADP-ribose). *Biochem. Biophys. Res. Commun.* 1985; 126:136-42.
- Smith S, Giriati I, Schmitt A, de Lange T. Tankyrase, a poly(ADP-ribose) polymerase at human telomeres. *Science* 1998c; 282:1484-7.

- Smith ST, Petruk Y, Sedkov Y, Cho E, Tillib S, Canaani E, Mazo A. Modulation of heat shock gene expression by the TAC1 chromatin-modifying complex. *Nat. Cell Biol.* 2004; 6:162-7.
- Sohn KJ, Stempak JM, Reid S, Shirwadkar S, Mason JB, Kim YI. The effect of dietary folate on genomic and p53-specific DNA methylation in rat colon. *Carcinogenesis* 2003;24:81-90.
- Soloaga A, Thomson S, Wiggin GR, Rampersaud N, Dyson MH, Hazzalin CA, Mahadevan LC, Arthur JS. MSK2 and MSK1 mediate the mitogen- and stress-induced phosphorylation of histone H3 and HMG-14. *EMBO J.* 2003;22:2788-97.
- Song F, Smith JF, Kimura MT, Morrow AD, Matsuyama T, Nagase H, Held WA. Association of tissue-specific differentially methylated regions (TDMs) with differential gene expression. *Proc. Natl. Acad. Sci. USA* 2005;102:3336-41.
- Southern EM. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 1975;98:503-17.
- Souza CP de, Osmani AH, Wu LP, Spotts JL, Osmani SA. Mitotic histone H3 phosphorylation by the NIMA kinase in *Aspergillus nidulans*. *Cell* 2000;102:293-302.
- Spadaccini R, Perrin H, Bottomley MJ, Ansieau S, Sattler M. Structure and function of the MYND domain. *J. Mol. Biol.* 2006;358:498-508.
- Spiegelman BM, Heinrich R. Biological control through regulated transcriptional coactivators. *Cell* 2004;119:157-67.
- Srinivasan M, Aalinkel R, Song F, Patel MS. Programming of islet functions in the progeny of hyperinsulinemic/obese rats. *Diabetes* 2003;52:984-90.
- St Clair D, Xu M, Wang P, Yu Y, Fang Y, Zhang F, Zheng X, Gu N, Feng G, Sham P, He L. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959-1961. *JAMA* 2005;557-62.
- Stadler WM, Margolin K, Ferber S, McCulloch W, Thompson JA. A phase II study of depsipeptide in refractory metastatic renal cell cancer. *Clin. Genitouterin Cancer* 2006;5:57-60.
- Stanley JS, Griffin JB, Zempleni J. Biotinylation of histones in human cells: effects of cell proliferation. *Eur. J. Biochem.* 2001;268:5424-9.
- Steele-Perkins G, Fang W, Yang XH, van Gele M, Carling T, Gu J, Buyse IM, Fletcher JA, Liu J, Bronson R, Chadwick RB, de la Chapelle A, Zhang X, Speleman F, Huang S. Tumor formation and inactivation of RIZ1, and Rb-binding member of a nuclear protein-methyltransferase superfamily. *Genes Dev.* 2001;15:2250-62.
- Steenman MJ, Rainier S, Dobry CJ, Grundy P, Horon IL, Feinberg AP. Loss of imprinting of IGF2 is linked to reduced expresión and abnormal methylation of H19 in Wilms' tumor. *Nature Genet.* 1994;7:433-9.
- Stein AD, Lumey LH. The relationship between maternal and offspring birth weights after maternal prenatal famine exposure: the Dutch famine birth cohort study. *Hum. Biol.* 2000; 72:641-54.
- Stern DE, Berger SL. Acetylation of histones and transcription-related factors. *Microbiol. Mol. Biol. Rev.* 2000; 64:435-59.
- Stern DE, Belotserkovskaya R, Berger SL. ALSA, a variant of yeast SAGA, containg truncated Spt7, which correlates with activated transcription. *Proc. Natl. Acad. Sci. USA* 2002;99:11622-7.
- Stewart DJ, Donehower RC, Eisenhauer EA, Wainman N, Shah AK, Bonfils C, MacLeod AR, Besterman JM, Reid GK. A phase I pharmacokinetic and pharmacodynamic study of the DNA methyltransferase 1 inhibitor MG98 administered twice weekly. *Ann. Oncol.* 2003; 14:766-74.
- Stewart RJC, Preece RF, Sheppard HG. Twelve generations of marginal protein deficiency. *Br. J. Nutr.* 1975; 33:233-53.
- Stocken LA. ADP-ribosylation of nuclear proteins. *Methods Cell Biol.* 1978;19:167-74.
- Stockert RJ, Paietta E, Racevskis J, Morell AG. Posttranscriptional regulation of the asialoglycoprotein receptor by cGMP. *J. Biol. Chem.* 1992;267:56-9.
- Strahl BD, Allis CD. The language of covalent histone modifications. *Nature* 2000; 403:41-5.

- Strahl BD, Briggs SD, Brame CJ, Caldwell JA, Koh SS, Ma H, Cook RG, Shabanowitz J, Hunt DF, Stallcup MR, Allis CD. Methylation of histone H4 at arginine 3 occurs in vivo and is mediated by the nuclear receptor coactivator. *Curr. Biol.* 2001;11:996-1000.
- Strathdee G, Brown R. Aberrant DNA methylation in cancer: potential clinical interventions. *Expert Rev. Mol. Med.* 2002;4:1-17.
- Strathdee G, MacKean MJ, Illand M, Brown R. A role for methylation of the hMLH1 promoter in loss of hMLH1 expression and drug resistance in ovarian cancer. *Oncogene* 1999;18:2335-41.
- Streit WR, Entcheva P. Biotin in microbes, the genes involved in its biosynthesis, its biochemical role and perspectives for biotechnological production. *Appl. Microbiol. Biotechnol.* 2003;61:21-31.
- Strelkov IS, Davie JR. Ser-10 phosphorylation of histone H3 and immediate early gene expression in oncogene-transformed mouse fibroblasts. *Cancer Res.* 2002;62:75-8.
- Sun XJ, Wei J, Wu XY, Hu M, Wang L, Wang HH, Zhang QH, Chen SJ, Huang QH, Chen Z. Identification and characterization of a novel human histone H3 lysine 36-specific methyltransferase. *J. Biol. Chem.* 2005;280:35261-71.
- Sun ZW, Allis CD. Ubiquitination of histone H2B regulates H3 methylation and gene silencing in yeast. *Nature* 2002;418:104-8.
- Swango KL, Wolf B. Conservation of biotinidase in mammals and identification of the putative biotinidase gene in *Drosophila melanogaster*. *Mol. Genet. Metab.* 2001;74:492-9.
- Tachibana M, Sugimoto K, Fukushima T, Shinkai Y. SET domain-containing protein, G9a, is a novel lysine-preferring mammalian histone methyltransferase with hyperactivity and specific selectivity to lysine 9 and 27 of histone H3. *J. Biol. Chem.* 2001;276:25309-17.
- Tajun-Arifin K, Teasdale R, Ravasi T, Hume DA, Mattick JS. Identification and analysis of chromodomain-containing proteins encoded in the mouse transcriptome. *Genome Res.* 2003; 13:1416-29.
- Takahashi M, Kamei Y, Ezaki O. Mest/Peg1 imprinted gene enlarges adipocytes and is a marker of adipocyte size. *Am. J. Physiol. Endocrinol. Metab.* 2005;288:E117-E124.
- Takahashi Y, Kahyo T, Toh EA, Yasuda H, Kikuchi Y. Yeast Ull1/Siz1 is a novel SUMO1/Smt3 ligase for septin components and functions as an adaptor between conjugating enzyme and substrates. *J. Biol. Chem.* 2001;276:48973-7.
- Takechi S, Nakayama T. Sas3 is a histone acetyltransferase and requires a zinc finger motif. *Biochem. Biophys. Res. Commun.* 1999;266:405-10.
- Takiguchi M, Achazanar WE, Qu W, Li G, Waalkes MP. Effects of cadmium on DNA-(cytosine-5) methyltransferase activity and DNA methylation status during cadmium-induced cellular transformation. *Exp. Cell Res.* 2003;286:355-65.
- Tamaru H, Selker EU. A histone methyltransferase controls DNA methylation in *Neurospora crassa*. *Nature* 2001;414:277-83.
- Tan J, Zhuang L, Jiang X, Yang KK, Karuturi KM, Yu Q. Apoptosis signal-regulating kinase 1 is a direct target of E2F1 and contributes to histone deacetylase inhibitor-induced apoptosis through positive feedback regulation of E2F1 apoptotic activity. *J. Biol. Chem.* 2006;281:10508-15.
- Tan K, Shaw AL, Madsen B, Jensen K, Taylor-Papadimitriou J, Freemont PS. Human PLU-1 has transcriptional repression properties and interacts with the developmental transcription factors BF-1 and PAX9. *J. Biol. Chem.* 2003;278:20507-13.
- Tanigawa Y, Tsuchiya M, Imai Y, Shimoyama M. ADP-ribosyltransferase from hen liver nuclei. Purification and characterization. *J. Biol. Chem.* 1984;259:2022-9.
- Tanny JC, Dowd GJ, Huang J, Hilz H, Moazed D. An enzymatic activity in the yeast Sir2 protein is essential for gene silencing. *Cell* 1999;99:735-45.
- Tanuma S, Kawashima K, Endo H. Acceptor proteins for (ADP-ribose)n in the HeLa S3 cell cycle. *J. Biochem. (Tokyo)* 1986;99:915-22.

- Tatematsu KI, Yamazaki T, Ishikawa F. MBD2-MBD3 complex binds to hemi-methylated DNA and forms a complex containing DNMT1 at the replication foci in late S phase. *Genes Cells* 2000;5:677-88.
- Tatham MH, Jaffray E, Vaughan OA, Desterro JM, Botting CH, Naismith JH, Hay RT. Polymeric chains of SUMO-2 and SUMO-3 are conjugated to protein substrates by SAE1/SAE2 and Ubc9. *J. Biol. Chem.* 2001;276:35368-74.
- Thakur N, Tiwari BK, Thomassin H, Pandey RR, Kanduri M, Göndör A, Grange T, Ohlsson R, Kanduri C. An antisense RNA regulates the bidirectional silencing property of the Kcnq1 imprinting control region. *Mol. Cell Biol.* 2004;24:7855-62.
- Thatcher KN, Peddada S, Yasui DH, LaSalle JM. Homologous paring of 15q11-13 imprinted domains in brain is developmentally regulated but deficient in Rett and autism samples. *Hum. Mol. Genet.* 2005;14:785-97.
- Thomas T, Voss AK. The diverse biological roles of MYST histone acetyltransferase family proteins. *Cell Cycle* 2007;6:696-704.
- Thomas T, Voss AK, Chowdhury K, Gruss P. Querkopf, a MYST family histone acetyltransferase, is required for normal cerebral cortex development. *Development* 2000;127:2537-48.
- Thomson S, Mahadevan LC, Clayton AL. MAP kinase-mediated signalling to nucleosomes and immediate-early gene induction. *Sem. Cell Dev. Biol.* 1999;10:205-14.
- Thorne AW, Sautiere P, Briand G, Crane-Robinson C. The structure of ubiquitinated histone H2B. *EMBO J.* 1987;5:1005-10.
- Torrens C, Brawley L, Dance CS, Itoh S, Poston L, Hanson MA. First evidence for transgenerational vascular programming in the rat protein restriction model. *J Physiol* 543;41P-42P.
- Toyota M, Ho C, Ahuja N, Jair KW, Li Q, Ohe-Toyota M, Baylin SB, Issa JP. Identification of differentially methylated sequences in colorectal cancer by methylated CpG island amplification. *Cancer Res.* 1999;59:2307-12.
- Trasler J, Deng L, Melnyk S, Pogribny I, Hiou-Tim F, Sibani S, Oakes C, Li E, James SJ, Rozen R. Impact of DNMT1 deficiency, with and without low folate diets, on tumor numbers and DNA methylation in Min mice. *Carcinogenesis* 2003;24:39-45.
- TREMEDICAPanace@ Asociación Internacional de Traductores y Redactores de Medicina y Ciencias Afines. Disponible EN [http://www.medtrad.org/info\\_general.html](http://www.medtrad.org/info_general.html). Acesado en: 20 de abril 2008.
- Tremolizzo L, Carboni G, Ruzicka WB, Mitchell CP, Sugaya I, Tueting P, Sharma R, Grayson DR, Costa E, Guidotti A. An epigenetic mouse model for molecular and behavioral neuropathologies related to schizophrenia vulnerability. *Proc. Natl. Acad. Sci. USA* 2002;99:17095-100.
- Trivedi CM, Luo Y, Yin Z, Zhang M, Zhu W, Wang T, Floss T, Goettlicher M, Noppinger PR, Wurst W, Ferrari VA, Abrams CS, Gruber PJ, Epstein JA. HDAC2 regulates the cardiac hypertrophic response by modulating Gsk3b activity. *Nat. Med.* 2007;13:324-31.
- Trujillo E, Davis C, Milner J. Nutrigenomics, proteomics, metabolomics and the practice of dietetics. *J. Am. Diet. Assoc.* 2006; 106:403-13.
- Tsujiuchi T, Tsutsumi M, Sasaki Y, Takama M, Konishi Y. Hypomethylation of CpG sites and c-myc gene overexpression in hepatocellular carcinomas, but not hyperplastic nodules, induced by a choline-deficient L-aminoacid-defined diet in rats. *Jpn. J. Cancer Res.* 1999;90:909-13.
- Tsukada Y, Fang J, Erdjument-Bromage H, Warren ME, Borchers CH, Tempst P, Zhang Y. Histone demethylation by a family of JmjC domain-containing proteins. *Nature* 2006;439:811-6.
- Turner BM. Histone acetylation and an epigenetic code. *Bioessays* 2000; 22:836-45.
- Turner BM. *Chromatin and Gene Regulation: Molecular Mechanisms in Epigenetics*. Birmingham, UK: Blackwell Science; 2001.
- Turner BM. Reading signals on the nucleosome with a new nomenclatura for modified histones. *Nat. Struct. Mol. Biol.* 2005; 12:110-2.
- Turner SD, Ricci AR, Petropoulos H, Genereaux J, Skerjanc IS, Brandl CJ. The E2 ubiquitin conjugase Rad6 is required for the Arar/Mcm1 repression of ARG1 transcription. *Mol. Cell Biol.* 2002; 22:4011-9.

- Usadel H, Brabender J, Danenberg KD, Jerónimo C, Harden S, Engles J, Danenberg PV, Yang S, Sidransky D. Quantitative adenomatous polyposis coli promoter methylation analysis in tumor tissue, serum, and plasma DNA of patients with lung cancer. *Cancer Res.* 2002;62:371-5.
- Ushiroyama T, Tanigawa Y, Tsuchiya M, Mishima K, Yoshimura Y, Shimoyama M. Amino acid sequence of histone H1 at the ADP-ribose-accepting site and ADP-ribose X histone-H1 adduct as an inhibitor of cyclic-ADM-dependent. *Eur. J. Biochem.* 1985;151:173-7.
- USNLM Genetics Home Reference. U.S. National Library of Medicine. Disponible en [http://ghr.nlm.nih.gov/glossary=genetic\\_heterogeneity](http://ghr.nlm.nih.gov/glossary=genetic_heterogeneity). Accesado el 19 de junio 2008a.
- USNLM Medline Plus Encyclopedia en español. Disponible en <http://www.nlm.nih.gov/medlineplus/spanish/encyclopedia.html>. Accesado el 10 de junio 2008b.
- Uthus EO. Estimation of safe and adequate daily intake for arsenic. En: Mertz W, Abernathy CO, Olin SS (eds). *Risk Assessment of Essential Elements*. Washington DC; 1994.
- Vadlamudi S, Kalhan SC, Patel MS. Persistence of metabolic consequences in the progeny of rats fed a HC formula in their early postnatal life. *Am. J. Physiol.* 1995; 269:E731-8.
- Varionostic 'Pyrosequencing strengths and applications.' Varionostic bioanalytical services. Disponible en [http://www.varionostic.de/fileadmin/downloads/pyroprinciple\\_descr.pdf](http://www.varionostic.de/fileadmin/downloads/pyroprinciple_descr.pdf). Accesado el 31 de mayo 2008.
- Vaziri H, Dessain SK, Ng Eaton E, Imai SI, Frye RA, Pandita TK, Guarante L, Weinberg RA. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* 2001;107:149-59.
- Vecchia C la, Negri E, Pelucchi C, Franceschi S. Dietary folate and colorectal cancer. *Int. J. Cancer* 2002;102:545-7.
- Vega RB, Matsuda K, Oh J, Barbosa AC, Yang X, Meadows E, McAnally J, Pomajzl C, Shelton JM, Richardson JA, Karsenty G, Olson EN. Histone deacetylase 4 controls chondrocyte hypertrophy during skeletogenesis. *Cell* 2004;119:555-66.
- Veldic M, Guidotti A, Maloku E, Davis JM, Costa E. In psychosis, cortical interneurons overexpress DNA-methyltransferase 1. *Proc. Natl. Acad. Sci. USA* 2005;102:2152-7.
- Verger A, Perdomo J, Crossley M. Modification with SUMO. A role in transcriptional regulation. *EMBO Rep.* 2003;4:137-42.
- Vickaryous N, Whitelaw E. The role of early embryonic environment on epigenotype and phenotype. *Reprod. Fertil. Dev.* 2005; 17:335-40.
- Vidali G, Boffa LC, Bradbury EM, Allfrey VG. Butyrate suppression of histone deacetylation leads to accumulation of multiacetylated forms of histones H3 and H4 and increased DNase I sensitivity of the associated DNA sequences. *Proc. Natl. Acad. Sci.* 1978;75:2239-43.
- Vijay-Kumar S, Bugg CE, Cook WJ. Structure of ubiquitina refined at 1.8 Å resolution. *J. Mol. Biol.* 1987;194:531-44.
- Vijg J, Suh Y. Ageing: chromatin unbound. *Nature* 2006;440:874-5.
- Villar-Garea A, Frage MF, Espada J, Esteller M. Procaine is a DNA-demethylating agent with growth-inhibitory effects in human cancer cells. *Cancer Res.* 2003;63:4984-9.
- Viré E, Brenner C, Deplus R, Blanchon L, Fraga M, Didelot C, Morey L, Van Eynde A, Bernard D, Vanderwinden JM, Bollen M, Esteller M, Di Croce L, de Launoit Y, Fuks F. The polycomb group protein EZH2 directly controls DNA methylation. *Nature* 2006;439:871-4.
- Visapää JP, Götte K, Benesova M, Li J, Homann N, Conradt C, Inoue H, Tisch M, Hörrmann K, Väkeväinen S, Salaspuro M, Seitz HK. Increased cancer risk in heavy drinkers with the alcohol dehydrogenase 1C\*1 allele, possibly due to salivary acetaldehyde. *Gut* 2004;53:871-6.
- Vliet J van, Oates NA, Whitelaw E. Epigenetic mechanisms in the context of complex diseases. *Cell Mol. Life Sci.* 2007;64:1531-8.
- Volkel P, Angrand PO. The control of histone lysine methylation in epigenetic regulation. *Biochimie* 2007;89:1-20.
- Waddington CH. Organisers and genes. Cambridge: Cambridge Univ. Press; 1940.

- Wade PA, Pruss D, Wolffe AP. Histone acetylation: Chromatin in action. *Trends Biochem. Sci.* 1997;22:128-32.
- Wade PA. Transcriptional control at regulatory checkpoints by histone deacetylases: molecular connections between cancer and chromatin. *Hum. Mol. Genet.* 2001;10:693-8.
- Wallwork JC, Duerre JA. Effect of zinc deficiency on methionine metabolism, methylation reactions and protein synthesis in isolated perfused rat liver. *J. Nutr.* 1985;115:252-62.
- Walsh CP, Bestor TH. Cytosine methylation and mammalian development. *Genes Dev. S.* 1999;13:26-34.
- Wang H, Huang ZQ, Xia L, Feng Q, Erdjument-Bromage H, Strahl BD, Briggs SD, Allis CD, Wong J, Tempst P, Zhang Y. Methylation of histone H4 at arginine 3 facilitates transcriptional activation by nuclear hormone receptor. *Science* 2001;293:853-7.
- Wang L, Brown JL, Cao R, Zhang Y, Kassis JA, Jones RS. Hierarchical recruitment of polycomb group silencing complexes. *Mol. Cell* 2004a;14:637-48.
- Wang L, Mizzen C, Ying C. Histone acetyltransferase activity is conserved between yeast and human GCN5 and is required for complementation of growth and transcriptional activation. *Mol. Cell Biol.* 1997;17:519-27.
- Wang XF, Qian DZ, Ren M, Kato Y, Wei Y, Zhang L, Fansler Z, Clark D, Nakanishi O, Pili R. Epigenetic modulation of retinoic acid receptor beta2 by the histone deacetylase inhibitor MS-275 in human renal carcinoma. *Clin. Cancer Res.* 2005;11:3535-42.
- Wang Y, Griffiths WJ, Jörnvall H, Agerberth B, Johansson J. Antibacterial peptides in stimulated human granulocytes: characterization of ubiquitinated histone H1A. *Eur. J. Biochem.* 2002;269:512-8.
- Wang Y, Wusocka J, Sayegh J, Lee YH, Perlin JR, Leonelli L, Sonbuchner LS, McDonald CH, Cook RG, Dou Y, Roeder RG, Clarke S, Stallcup MR, Allis CD, Coonrod SA. Human PAD4 regulates histone arginine methylation levels via demethylimation. *Science* 2004b;306:279-83.
- Warnecke PM, Stirzaker C, Melki JR, Millar DS, Paul CL, Clark SJ. Optimizing annealing temperature overcomes bias in bisulfite PCR methylation analysis. *Nucleic Acid. Res.* 1997;25:4422-6.
- Waterland RA. Assessing the effects of high methionine intake on DNA methylation. *J. Nutr.* 2006;136:1706S-1710S.
- Waterland RA, Garza C. Potential mechanisms of metabolic imprinting that lead to chronic disease. *Am. J. Clin. Nutr.* 1999;69:179-97.
- Waterland RA, Jirtle RL. Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol. Cell Biol.* 2003;23:5293-300.
- Waterland RA, Jirtle RL. Early nutrition, epigenetic changes at transposons and imprinted genes, and enhanced susceptibility to adult chronic diseases. *Nutrition* 2004;20:63-8.
- Waterland RA, Lin JR, Smith CA, Jirtle RL. Postweaning diet affects genomic imprinting at the insulin-like growth factor 2 (Igf2) locus. *Hum. Mol. Genet.* 2006;15:705-16.
- Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M, Meaney MJ. Epigenetic programming by maternal behavior. *Nat. Neurosci.* 2004;7:847-54.
- Weber M, Davies JJ, Witting D, Oakeley EJ, Haase M, Lam WL, Schübeler D. Chromosome-wide and promoter-specific analyses identify sites of differential DNA methylation in normal and transformed human cells. *Nat. Genet.* 2005;37:853-62.
- Weisenberger DJ, Campan M, Long TI, Kim M, Woods C, Fiala E, Ehrlich M, Laird PW. Analysis of repetitive element DNA methylation by MethylLight. *Nucleic Acids Res.* 2005;33:6823-36.
- Weksberg R, Smith AC, Squire J, Sadowski P. Beckwith-Wiedemann syndrome demonstrates a role for epigenetic control of normal development. *Hum. Mol. Genet.* 2003;12:R61-R68.
- West MH, Bonner WM. Histone 2B can be modified by the attachment of ubiquitina. *Nucleic Acids Res.* 1989;8:4671-80.
- Whetstone JR, Nottke A, Lan F, Huarte M, Smolikov S, Chen Z, Spooner E, Li E, Zhang G, Colaiacovo M, Shi Y. Reversal of histone lysine trimethylation by the JMJD2 family of histone demethylases. *Cell* 2006;125:467-81.

- Wielckens K, Bredehorst R, Adamietz P, Hilz H. Mono-ADP-ribosylation and poly-ADP-ribosylation of proteins in normal and malignant tissues. *Adv. Enzyme Regul.* 1982;20:23-37.
- Wijermans P, Lubbert M, Verhoef G, Bosly A, Ravoet C, Andre M, Ferrant A. Low dose 5-aza-2'-deoxycytidine, a DNA hypomethylating agent, for the treatment of high-risk myelodysplastic syndrome: a multicenter phase II study in elderly patients. *J. Clin. Oncol.* 2000;18:956-62.
- Wikipedia. Ubiquitin. Disponible en: <http://en.wikipedia.org/wiki/Ubiquitin>. Accesado el 16 de mayo 2008.
- Wilhelm J, Bayerlein K, Hilemacher T, Reulbach U, Frieling H, Kromolan B, Degner D, Kornhuber J, Bleich S. Short-term cognition deficits during early alcohol withdrawal are associated with elevated plasma homocysteine levels in patients with alcoholism. *J. Neural Transm.* 2005;113:357-63.
- Wilkinson KD. Ubiquitination and deubiquitination: Targeting of proteins for degradation by the proteasome. *Semin. Cell Dev. Biol.* 2000;11:141-8.
- Williams RR, Azuara V, Perry P, Sauer S, Dvorkina M, Jørgensen H, Roix J, McQueen P, Misteli T, Merkenschlager M, Fisher AG. Neural induction promotes large-scale chromatin reorganisation o the Mash1 locus. *J. Cell Sci.* 2006;119:132-40.
- Wilson JR, Jing C, Walker PA, Martin SR, Howell SA, Blackburn GM, Gamblin SJ, Xiao B. Crystal structure and functional analysis of the histone methyltransferase SET7/9. *Cell* 2002;111:105-15.
- Wilson VL, Jones PA. DNA methylation decreases in aging but not in immortal cells. *Science* 1983;220:1055-7.
- Wolf B. *The Metabolic and Molecular Bases of Inherited Disease*. Nueva York: McGraw-Hill; 2001.
- Wolfe J. Introduction to human genetics. Glossary of terms used in molecular genetics. Disponible en: <http://www.ucl.ac.uk/~ucbhjow/b241/glossary.html>. Accesado el 19 de junio 2008.
- Wolff GL, Kodell RL, Moore SR. Maternal epigenetics and methyl supplements affect *agouti* gene expression in A<sup>y/a</sup> mice. *FASEB J.* 1998;12:949-57.
- Wong IH, Lo YM, Yeo W, Lau WY, Johnson PJ. Frequent p15 promoter methylation in tumor and peripheral blood from hepatocellular carcinoma patients. *Clin. Cancer Res.* 2000;6:3516-21.
- Wray GA. The evolutionary significance of cis-regulatory mutations. *Nat. Rev. Genet.* 2007; 8:206-16.
- Wu G, Bazer FW, Cudd TA, Meininger CJ, Spencer TE. Maternal nutrition and fetal development. *J. Nutr.* 2004;134:2169-72.
- Wu Q, Suzuki M. Parental obesity and overweight affect the body-fat accumulation in the offspring: the possible effect of a high-fat diet through epigenetic inheritance. *Obes. Rev.* 2006;7:201-8.
- Wu RS, Khon KW, Bonner WM. Metabolism of ubiquitinated histones. *J. Biol. Chem.* 1981;256:5916-20.
- Wu Y, Guo SW. Inhibition of proliferation of endometrial stromal cells by trichostatin A, RU486, CDB-2914, N-acetylcysteine, and ICI 182780. *Gynecol. Obstet. Invest.* 2006; 62:193-205.
- Wyngaert I van den, de Vries W, Kremer A, Neefs J, Verhasselt P, Luyten WH, Kass SU. Cloning and characterization of human histone deacetylase 8. *FEBS Lett.* 2000;478:77-83.
- Wysocka J, Swigut T, Milne TA, Dou Y, Zhang X, Burlingame AL, Roeder RG, Brivanlou AH, Allis CD. WDR5 associates with histone H3 methylated at K4 and is essential for H3-K4 methylation and vertebrate development. *Cell* 2005;121:859-72.
- Wysocka J, Swigut T, Xiao H, Milne TA, Kwon SY, Landry J, Kauer M, Tackett AJ, Chait BT, Badenhorst P, Wu C, Allis CD. A PHD finger of NURF couples histone H3 lysine 4 trimethylatiion with chromatin remodelling. *Nature* 2006;442:86-90.
- Xiao B, Jing C, Wilson JR, Walker PA, Vasishth N, Kelly G, Howell S, Taylor IA, Blackburn GM, Gamblin SJ. Structure and catalytic mechanism of the human histone methyltransferase SET7/9. *Nature* 2003a; 421:652-6.
- Xiao T, Hall H, Kizer KO, Shibata Y, Hall MC, Borchers CH, Strahl BD. Phosphorylation of RNA polymerase II CTD regulates H3 methylation in yeast. *Genes Dev.* 2003b; 17:654-63.
- Xiao T, Kao CF, Krogan NJ, Sun ZW, Greenblatt JF, Osley MA, Strahl BD. Histone H2B ubiquitylation is associated with elongating RNA polymerase II. *Mol. Cell Biol.* 2005; 25:637-51.
- Xie S, Wang Z, Okano M, Nogami M, Li Y, He WW, Okumura K, Li E. Cloning, expression and chromosome locations of the human DNMT3 gene family. *Gene* 1999;236:87-95.

- Xin Z, Tachibana M, Guggiari M, Heard E, Shinkai Y, Wagstaff J. Role of histone methyltransferase G9a in CpG methylation of the Prader Willi syndrome imprinting center. *J. Biol. Chem.* 2003;278:14996-15000.
- Xiong Z, Laird PW. COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res.* 1997;25:2532-4.
- Xu C, Henry MF. Nuclear export of hnRNP Hrp1p and nuclear export of hnRNP NpI3p are linked and influenced by the methylation state of NpI3p. *Mol. Cell Biol.* 2004;24:10742-56.
- Xu W, Chen H, Du K, Asahara H, Tini M, Emerson BM, Montminy M, Evans RM. A transcriptional switch mediated by cofactor methylation. *Science* 2001;294:2507-11.
- Xu W, Edmondson DG, Roth SY. Mammalian GCN5 and P/CAF acetyltransferases have homologous amino-terminal domains important for recognition of nucleosomal substrates. *Mol. Cell Biol.* 1998; 18:5655-69.
- Xu WS, Parmigiani R, Marks PA. Histone deacetylase inhibitors: molecular mechanism of action. *Oncogene* 2007;26:5541-52.
- Yamamoto T, Horikoshi M. Novel substrate specificity of the histone acetyltransferase activity of HIV-1-Tat interactive protein Tip60. *J. Biol. Chem.* 1997;272:30595-8.
- Yamane K, Toumazou C, Tsukada Y, Erdjument-Bromage H, Tempst P, Wong J, Zhang Y. JHDM2A, a JmjC-containing H3K9 demethylase, facilitates transcription activation by the androgen receptor. *Cell* 2006;125:483-95.
- Yan PS, Chen CM, Shi H, Rahmatpanah F, Wei SH, Huang TH. Applications of CpG island microarrays for high-throughput analysis of DNA methylation. *J. Nutr.* 2002;132:2430S-2434S.
- Yanagisawa J, Kitawa H, Yanagida M, Wada O, Ogawa S, Nakagomi M, Oishi H, Yamamoto Y, Nagasawa H, McMahon SB, Cole MD, Tora L, Takahashi N, Kato S. Nuclear receptor function requires a TFTC-type histone acetyl transferase complex. *Mol. Cell* 2002;9:553-62.
- Yang AS, Doshi KD, Choi SW, Mason JB, Mannari RK, Gharybian V, Luna R, Rashid A, Shen L, Estecio MR, Kantarjian HM, Garcia-Manero G, Issa JP. DNA methylation changes after 5-aza-2'-deoxycytidine therapy in patients with leukemia. *Cancer Res.* 2006;66:5495-5503.
- Yang AS, Estecio MR, Doshi K, Kondo Y, Tajara EH, Issa JP. A simple method for estimating global DNA methylation using bisulfite PCR of repetitive DNA elements. *Nucleic Acids Res.* 2004;32:E38.
- Yang L, Xia L, Wu DY, Wang H, Chansky HA, Schubach WH, Hickstein DD, Zhang Y. Molecular cloning of ESET, a novel histone H3-specific methyltransferase that interacts with ERG. *Oncogene* 2002a; 21:148-52.
- Yang SH, Jaffray E, Hay RT, Sharrocks AD. Dynamic interplay of the SUMO and ERK pathways in regulating Elk-1 transcriptional activity. *Mol. Cell* 2003;12:63-74.
- Yang WM, Tsai SC, Wen YD, Tejer G, Seto E. Functional domains of histone deacetylase-3. *J. Biol. Chem.* 2002b;277:9447-54.
- Yang XJ. The diverse superfamily of lysine acetyltransferases and their roles in leukaemia and other diseases. *Nucleic Acids Res.* 2004; 32:959-76.
- Yang XJ, Ogryzko VV, Nishikawa J, Howard BH, Najatani Y. A p300/CBP-associated factor that competes with the adenoviral oncoprotein E1A. *Nature* 1996;382:319-24.
- Yatabe Y, Mitsudomi T, Takahashi T. Maspin expression in normal lung and non-small-cell lung cancers: cellular property-associated expression under the control of promoter DNA methylation. *Oncogene* 2004;23:4041-9.
- Ye L, Dinkova-Kostova AT, Wade KL, Zhang Y, Shapiro TA, Talalay P. Quantitative determination of dithiocarbamates in human plasma, serum erythrocytes and urine: pharmacokinetics of broccoli sprout isothiocyanates in humans. *Drug Metab. Rev.* 2002;316:43-53.
- Yen R-WC, Vertino PM, Nelkin BD, Yu JJ, el-Deiry W, Cumaraswamy A, Lennon GG, Trask BJ, Celano P, Baylin SB. Isolation and characterization of the cDNA encoding human DNA methyltransferases. *Nucleic Acids Res.* 1992;20:2287-91.
- Yeung F, Hoberg JE, Ramsey CS, Keller MD, Jones DR, Frye RA, Mayo MW. Modulation of NF- $\kappa$ B-dependent transcripción and cell survival by the SIRT1 deacetylase. *EMBO J.* 2004;23:2369-80.

- Yin H, Chen ZX, Cen JN, Wang W, Duan WM, Yao L. 2005. Mechanism of differentiation and apoptosis in leukemia cells induced by tributyrin. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 2005;13:774–77
- Ying AK, Hassanain HH, Roos CM, Smiraglia DJ, Issa JPJ, Michler RE, Caligiuri M, Plass, Goldschmidt-Clermont PJ. Methylation of the estrogen receptor-alpha gene promoter is selectively increased in proliferating human aortic smooth muscle cells. *Cardiovasc. Res.* 2000;46:172-9.
- Yoder JA, Soman NS, Verdine HL, Bestor TH. DNA (cytosine-5)-methyltransferases in mouse cells and tissues: studies with a mechanism-based probe. *J. Mol. Biol.* 1997;270:385-95.
- Yoder JA, Bestor TH. A candidate mammalian DNA methyltransferase related to pmt1p of fission yeast. *Hum. Mol. Genet.* 1998;7:279-84.
- Yokochi T, Robertson KD. Preferential methylation of unmethylated DNA by mammalian de novo DNA methyltransferase Dnmt3a. *J. Biol. Chem.* 2002;277:11735-45.
- Yokomori N, Tawata M, Anaya T. DNA methylation modulates mouse leptin promoter activity during the differentiation of 3T3-L1 cells. *Diabetologia* 2002;45:140-8.
- Yokoyama A, Wang Z, Wysocka J, Sanyal M, Aufiero DJ, Kitabayashi I, Herr W, Cleary ML. Leukemia proto-oncoprotein MLL forms a SET1-like histone methyltransferase complex with menin to regulate Hox gene expression. *Mol. Cell Biol.* 2004;24:5639-49.
- Yoo CB, Cheng JC, Jones PA. Zebularine: a new drug for epigenetic therapy. *Biochem. Soc. Trans.* 2004;32:910-2.
- Yoo CB, Jones PA. Epigenetic therapy of cancer: past, present and future. *Nat. Rev. Drug Discov.* 2006;5:37-50.
- Yoshida M, Kijima M, Akita M, Beppu T. Potent and specific inhibition of mammalian histone deacetylase both in vivo and in vitro by trichostatin A. *J. Biol. Chem.* 1990;265:17174-9.
- Yoshida M, Furumai R, Nishiyama M, Komatsu Y, Nishino N, Horinouchi S. Histone deacetylase as a new target for cancer chemotherapy. *Cancer Chemother. Pharmacol.* 2001;48:S20-S26.
- Yu MC, Bachand F, McBride AE, Komili S, Casolari JM, Silver PA. Arginine methyltransferase affects interactions and recruitment of mRNA processing and export factors. *Genes Dev.* 2004;18:2024-35.
- Yu M, Schreek S, Cerni C, Schamberger C, Lesniewicz C, Poreba K, Vervoorts E, Walsemann J, Grotzinger G, Kremmer J, Mehraein Y, Mertsching, J, Kraft R, Austen M, Luscher-Firzlaff J, Luscher B. PARP-10, a novel Myc-interacting protein with poly(ADP-ribose) polymerase activity, inhibits transformation. *Oncogene* 2005;24:1982-93.f
- Zahradka O, Ebisuzaki K. A shuttle mechanism for DNA-protein interactions. The regulation of poly(ADP-ribose) polymerase. *Eur. J. Biochem.* 1982;127:579-85.
- Zambrano E, Matínez-Samoya PM, Bautista CJ, Deás M, Guillén L, Rodríguez-González GL, Guzmán C, Larrea, F, Nathanielsz, PW. Sex differences in transgenerational alterations of growth and metabolism in progeny (F2) of female offspring (F1) of rats fed a low protein diet during pregnancy and lactation. *J. Physiol.* 2005a; 566:225-36.
- Zambrano P, Segura-Pacheco B, Perez-Cardenas E, Cetina L, Revilla-Vazquez A, Taja-Chayeb L, Chavez-Blanco A, Angeles E, Cabrera G, Sandoval K, Trejo-Becerril C, Chanona-Vilchis J, Duenas-González A. A phase I study of hydralazine to demethylate and reactivate the expresión of tumor supresor genes. *BMC Cancer* 2005b; 5:44.
- Zapisek WF, Cronin GM, Lyn-Cook BD, Poirier LA. The onset of oncogene hypomethylation in the livers of rats fed methyl-deficient, aminoacid-defined diets. *Carcinogenesis* 1992;13:1869-72.s
- Zempleni J. Uptake, localization, and noncarboxylase roles of biotin. *Annu. Rev. Nutr.* 2005;25 :175-96.
- Zeng L, Zhou MM. Bromodomain: an acetyl-lysine binding domain[abstract]. *FEBS Lett* 2002; 513:124-8. <http://www.ncbi.nlm.nih.gov/pubmed/11911891>. Accesado el 7 de mayo 2008.
- Zeschtnigk M, Bohringer S, Price EA, Onadim Z, Masshöfer L, Lohmann DR. A novel real-time PCR assay for quantitative analysis of methylated alleles (QAMA): analysis of retinoblastoma locus. *Nucleic Acids Res.* 2004;32:E125.

- Zhang CL, McKinsey TA, Chang S, Antos CL, Hill JA, Olson EN. Class II histone deacetylases act as signal-responsive repressors of cardiac hypertrophy. *Cell* 2002a; 110:479-88.
- Zhang D, Yoon HG, Wong J. JMJD2A is a novel N-CoR-interacting protein and is involved in repression of the human transcription factor Achaete Scute-like homologue 2 (ASCL2/Hash2). *Mol. Cell Biol.* 2005; 25:6404-14.
- Zhang X, Zhou L, Cheng X. Crystal structure of the conserved core of protein arginine methyltransferase PRMT3. *EMBO J.* 2000;19:3509-19.
- Zhang X, Tamaru H, Khan SI, Horton JR, Keefe LJ, Selker EU, Cheng X. Structure of Neurospora SET domain protein DIM-5, a histone H3 lysine methyltransferase. *Cell* 2002b;111:117-127.
- Zhang X, Cheng X. Structure of predominant protein arginine methyltransferase PRMT1 and analysis of its binding to substrate peptides. *Structure* 2003a;11:509-20.
- Zhang X, Yang Z, Khan SI, Horton JR, Tamaru H, Selker EU, Cheng X. Structural basis for the product specificity of histone methyltransferases. *Mol Cell.* 2003b;12:177-85.
- Zhang Y. Transcriptional regulation by histone ubiquitination and deubiquitination. *Genes Dev.* 2003;17:2733-40.
- Zhang Y, Reinberg D. Transcription regulation by histone methylation: interplay between different covalent modifications of the core histone tails. *Genes Dev.* 2001;15:2343-60.
- Zhong S, Salomoni P, Pandolfi PP. The transcriptional role of PML and the nuclear body. *Nature Cell Biol.* 2000a;2:E85-E90.
- Zhong S, Salomoni P, Ronchetti S, Guo A, Ruggero D, Pandolfi PP. Promyelocytic leukaemia (PML) and Daxx participate in a novel nuclear pathway for apoptosis. *J. Exp. Med.* 2000b;191:631-9.
- Zhou L, Cheng X, Connolly BA, Dickman MJ, Hurd PJ, Hornby DP. Zebularine: a novel DNA methylation inhibitor that forms a covalent complex with DNA methyltransferases. *J. Mol. Biol.* 2002;321:591-9.
- Zink D, Amaral MD, Englmann A, Lang S, Clarke LA, Rudolph C, Alt F, Luther K, Braz C, Sadoni N, Rosenecker J, Schindelhauer D. Transcription-dependent spatial arrangements of CFTR and adjacent genes in human cell nuclei. *J. Cell. Biol.* 2004;166:815-25.
- Zinner R, Albinez H, Walter J, Peters AH, Cremer T, Cremer M. Histone lysine methylation patterns in human cell types are arranged in distinct three-dimensional nuclear zones. *Histochem. Cell Biol.* 2006;125:3-19.
- Zinner R, Teller K, Versteeg R, Cremer T, Cremer M. Biochemistry meets nuclear architecture: multicolor immuno-FISH for co-localization analysis of chromosome segments and differentially expressed gene loci with various histone methylations. *Adv. Enzyme Regul.* 2007;47:223-41.
- Zou HZ, Yu BM, Wang ZW, Sun JY, Cang H, Gao F, Li DH, Zhao R, Feng GG, Yi J. Detection of aberrant p16 methylation in the serum of colorectal cancer patients. *Clin. Cancer Res.* 2002;8:188-91.