



ANEXO 2

Resultados SPSS

Descriptives

Notes

Output Created		12-MAR-2008 10:23:53
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User defined missing values are treated as missing. All non-missing data are used.
Syntax		DESCRIPTIVES VARIABLES=pleasure arousal control approach avoidance satisfaction /STATISTICS=MEAN STDDEV MIN MAX .
Resources	Elapsed Time	0:00:00.09

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
pleasure	121	1.00	4.20	1.9058	.67334
arousal	122	1.00	4.50	2.7404	.66806
control	122	1.00	5.00	2.2541	.82571
approach	122	1.33	5.00	2.9699	.64247
avoidance	122	1.00	3.33	2.1284	.56764
satisfaction	122	1.25	4.75	3.0943	.59814
Valid N (listwise)	121				

Regression



Notes

Output Created	12-MAR-2008 10:26:22	
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User-defined missing values are treated as missing. Statistics are based on cases with no missing values for any variable used.
Syntax	REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT approach /METHOD=ENTER pleasure arousal control .	
Resources	Elapsed Time	0:00:00.27
	Memory Required Additional Memory Required for Residual Plots	2980 bytes 0 bytes

Variables Entered/Removed(b)

Model	Variables Entered	Variables Removed	Method
1	control, arousal, pleasure(a)	.	Enter

a All requested variables entered.

b Dependent Variable: approach

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.237(a)	.056	.032	.62988

a Predictors: (Constant), control, arousal, pleasure

ANOVA(b)



Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	2.775	3	.925	2.331	.078(a)
	Residual	46.419	117	.397		
	Total	49.194	120			

a Predictors: (Constant), control, arousal, pleasure

b Dependent Variable: approach

Coefficients(a)

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	3.111	.261		11.898	.000
	pleasure	-.250	.096	-.263	-2.614	.010
	arousal	.081	.096	.085	.846	.399
	control	.047	.081	.061	.586	.559

a Dependent Variable: approach

Regression

Notes

Output Created		12-MAR-2008 10:27:03
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any variable used.
Syntax		REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT avoidance /METHOD=ENTER pleasure arousal control .
Resources	Elapsed Time	0:00:00.04
	Memory Required	2980 bytes
	Additional	0 bytes



Memory Required
for Residual Plots

Variables Entered/Removed(b)

Model	Variables Entered	Variables Removed	Method
1	control, arousal, pleasure(a)	.	Enter

a All requested variables entered.
b Dependent Variable: avoidance

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.166(a)	.027	.002	.56463

a Predictors: (Constant), control, arousal, pleasure

ANOVA(b)

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	1.051	3	.350	1.099	.353(a)
	Residual	37.300	117	.319		
	Total	38.351	120			

a Predictors: (Constant), control, arousal, pleasure
b Dependent Variable: avoidance

Coefficients(a)

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	1.877	.234		8.007	.000
	pleasure	.144	.086	.172	1.683	.095
	arousal	.010	.086	.012	.120	.905
	control	-.020	.072	-.029	-.276	.783

a Dependent Variable: avoidance

Regression

Notes

Output Created	12-MAR-2008 10:27:45
Comments	



Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User-defined missing values are treated as missing. Statistics are based on cases with no missing values for any variable used.
Syntax		REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT satisfaction /METHOD=ENTER pleasure arousal control .
Resources	Elapsed Time	0:00:00.05
	Memory Required Additional Memory Required for Residual Plots	2980 bytes 0 bytes

Variables Entered/Removed(b)

Model	Variables Entered	Variables Removed	Method
1	control, arousal, pleasure(a)	.	Enter

a All requested variables entered.

b Dependent Variable: satisfaction

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.364(a)	.132	.110	.56371

a Predictors: (Constant), control, arousal, pleasure

ANOVA(b)

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	5.678	3	1.893	5.956	.001(a)
	Residual	37.179	117	.318		



Total	42.857	120			
-------	--------	-----	--	--	--

a Predictors: (Constant), control, arousal, pleasure

b Dependent Variable: satisfaction

Coefficients(a)

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	3.880	.234		16.582	.000
	pleasure	-.242	.086	-.273	-2.826	.006
	arousal	-.215	.086	-.241	-2.503	.014
	control	.115	.072	.159	1.587	.115

a Dependent Variable: satisfaction

Regression

Notes

Output Created		12-MAR-2008 10:28:14
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User-defined missing values are treated as missing. Statistics are based on cases with no missing values for any variable used.
Syntax		REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT satisfaction /METHOD=ENTER pleasure arousal .
Resources	Elapsed Time	0:00:00.03
	Memory Required Additional	2684 bytes
	Memory Required for Residual Plots	0 bytes

Variables Entered/Removed(b)



Model	Variables Entered	Variables Removed	Method
1	arousal, pleasure(a)	.	Enter

a All requested variables entered.
 b Dependent Variable: satisfaction

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.337(a)	.114	.099	.56732

a Predictors: (Constant), arousal, pleasure

ANOVA(b)

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	4.878	2	2.439	7.579	.001(a)
	Residual	37.979	118	.322		
	Total	42.857	120			

a Predictors: (Constant), arousal, pleasure
 b Dependent Variable: satisfaction

Coefficients(a)

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	3.934	.233		16.879	.000
	pleasure	-.197	.081	-.222	-2.425	.017
	arousal	-.171	.082	-.192	-2.093	.038

a Dependent Variable: satisfaction

Univariate Analysis of Variance

Notes

Output Created	12-MAR-2008 10:41:24		
Comments			
Input	Filter	<none>	
	Weight	<none>	
	Split File	<none>	
	N of Rows in Working Data File	129	
Missing Value Handling	Definition of Missing Cases Used	User-defined missing values are treated as missing. Statistics are based on all cases with valid data for all variables in the	



Syntax		model.
		UNIANOVA approach BY F20 /METHOD = SSTYPE(3) /INTERCEPT = INCLUDE /POSTHOC = F20 (SCHEFFE) /EMMEANS = TABLES(OVERALL)
Resources	Elapsed Time	
		0:00:00.07

Between-Subjects Factors

	N
F20 1.00	48
2.00	20
3.00	19
4.00	3
5.00	3
6.00	29

Tests of Between-Subjects Effects

Dependent Variable: approach

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	2.172(a)	5	.434	1.055	.389
Intercept	356.791	1	356.791	866.339	.000
F20	2.172	5	.434	1.055	.389
Error	47.773	116	.412		
Total	1126.056	122			
Corrected Total	49.945	121			

a R Squared = .043 (Adjusted R Squared = .002)

Estimated Marginal Means

Grand Mean

Dependent Variable: approach

Mean	Std. Error	95% Confidence Interval	
		Lower Bound	Upper Bound
2.859	.097	2.666	3.051

Post Hoc Tests

F20



Multiple Comparisons

Dependent Variable: approach
Scheffe

(I) F20	(J) F20	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
1.00	2.00	.0069	.17080	1.000	-.5713	.5852
	3.00	.2065	.17394	.922	-.3824	.7954
	4.00	.0486	.38191	1.000	-1.2444	1.3416
	5.00	.7708	.38191	.542	-.5222	2.0639
	6.00	.1061	.15093	.992	-.4049	.6171
2.00	1.00	-.0069	.17080	1.000	-.5852	.5713
	3.00	.1996	.20559	.966	-.4965	.8956
	4.00	.0417	.39733	1.000	-1.3035	1.3869
	5.00	.7639	.39733	.596	-.5813	2.1091
	6.00	.0991	.18653	.998	-.5324	.7307
3.00	1.00	-.2065	.17394	.922	-.7954	.3824
	2.00	-.1996	.20559	.966	-.8956	.4965
	4.00	-.1579	.39869	.999	-1.5077	1.1919
	5.00	.5643	.39869	.847	-.7855	1.9142
	6.00	-.1004	.18941	.998	-.7417	.5409
4.00	1.00	-.0486	.38191	1.000	-1.3416	1.2444
	2.00	-.0417	.39733	1.000	-1.3869	1.3035
	3.00	.1579	.39869	.999	-1.1919	1.5077
	5.00	.7222	.52398	.862	-1.0518	2.4962
	6.00	.0575	.38920	1.000	-1.2602	1.3752
5.00	1.00	-.7708	.38191	.542	-2.0639	.5222
	2.00	-.7639	.39733	.596	-2.1091	.5813
	3.00	-.5643	.39869	.847	-1.9142	.7855
	4.00	-.7222	.52398	.862	-2.4962	1.0518
	6.00	-.6648	.38920	.713	-1.9825	.6530
6.00	1.00	-.1061	.15093	.992	-.6171	.4049
	2.00	-.0991	.18653	.998	-.7307	.5324
	3.00	.1004	.18941	.998	-.5409	.7417
	4.00	-.0575	.38920	1.000	-1.3752	1.2602
	5.00	.6648	.38920	.713	-.6530	1.9825

Based on observed means.

Homogeneous Subsets

approach

Scheffe

F20	N	Subset
		1
5.00	3	2.2778
3.00	19	2.8421
6.00	29	2.9425
4.00	3	3.0000
2.00	20	3.0417
1.00	48	3.0486



Sig.	.392
------	------

Means for groups in homogeneous subsets are displayed. Based on Type III Sum of Squares The error term is Mean Square(Error) = .412.

a Uses Harmonic Mean Sample Size = 7.276.

b The group sizes are unequal. The harmonic mean of the group sizes is used. Type I error levels are not guaranteed.

c Alpha = .050.

T-Test

Notes

Output Created	12-MAR-2008 10:51:47	
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User defined missing values are treated as missing. Statistics for each analysis are based on the cases with no missing or out-of-range data for any variable in the analysis.
Syntax	T-TEST GROUPS = F21(1 2) /MISSING = ANALYSIS /VARIABLES = satisfaction /CRITERIA = CI(.95) .	
Resources	Elapsed Time	0:00:00.04

Group Statistics

	F21	N	Mean	Std. Deviation	Std. Error Mean
satisfacti	1.00	117	3.0897	.58119	.05373
on	2.00	5	3.2000	1.00623	.45000

Independent Samples Test



+		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	Df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
satisfaction	Equal variances assumed	1.458	.230	-.402	120	.688	-.11026	.27411	-.65297	.43245
	Equal variances not assumed			-.243	4.115	.819	-.11026	.45320	-1.35480	1.13429

T-Test

Notes

Output Created	12-MAR-2008 10:54:01	
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	129
Missing Value Handling	Definition of Missing Cases Used	User defined missing values are treated as missing. Statistics for each analysis are based on the cases with no missing or out-of-range data for any variable in the analysis.
Syntax		T-TEST GROUPS = F2(1 2) /MISSING = ANALYSIS /VARIABLES = satisfaction /CRITERIA = CI(.95) .
Resources	Elapsed Time	0:00:00.04

Group Statistics

	F2	N	Mean	Std. Deviation	Std. Error Mean
satisfaction	1.00	49	3.2092	.64821	.09260
	2.00	73	3.0171	.55329	.06476



Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	Df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
satisfaction	Equal variances assumed	4.078	.046	1.753	120	.082	.19206	.10953	-.02480	.40892
	Equal variances not assumed			1.700	91.792	.093	.19206	.11300	-.03237	.41649